

# Wild salmon are getting smaller and it might be linked to aquaculture

February 25 2022

---



Atlantic salmon in the River Teno in Northern Finland. Credit: Panu Orell / Natural Research Institute Finland

Researchers from Finland used genetic methods to pinpoint how a fishery for an aquaculture fish food source and changes in salmon fishing may be linked to changes in the size of wild salmon.

The study, published recently in journal *Science*, showed the shrinking

size of Atlantic salmon in the River Teno in Northern Finland might not be due to directly fishing salmon. Rather, the impacts could stem from an indirect effect: the commercial fishery of one of wild salmon's favorite foods in the ocean: a small omega-3 rich [fish](#) called capelin.

This indirect effect identified in the study puts the spotlight on salmon aquaculture. Some of the capelin fishery catch is used as fishmeal for salmon aquaculture food, suggesting strong harvest and declining capelin abundance can be an indirect way salmon aquaculture could influence wild salmon populations.

"The [aquaculture industry](#) has made important progress in finding alternative protein sources for aquaculture fish feed, and our study suggests that these efforts have not been in vain, as it seems capelin harvest may affect wild salmon populations. Globally, 18 million tons of wild fish such as capelin are harvested annually for domestic animal feed, so there is still work to do to further reduce the effects of aquaculture on wild fish populations," notes professor Craig Primmer at the University of Helsinki.

## **The study links environmental and human influences with evolutionary changes**

"Our earlier research had shown that the age at which salmon were maturing in this river was getting younger, and consequently also the size of salmon that are spawning was getting smaller, showing "evolution in action." Important for demonstrating rapid evolution, there were also changes in their DNA at a gene known to be linked with maturation size and age," explains Primmer.

"This previous research couldn't tell us what environmental or human influences might be linked with the evolutionary changes. To understand

this, we needed to link the yearly changes in the salmon DNA variation with annual changes in environmental and human-linked factors," Dr. Yann Czorlich, the first author of the research continues. "We gathered literally millions of data-points about factors, including yearly water temperature, [salmon fishing](#) effort, and commercial fisheries catches of the fish salmon eat in the ocean, and compared them with our data on DNA changes in our 40-year time series"

## **Changes in salmon fishing affect the size of salmon**

In addition to the indirect effect of capelin harvest, the team also identified a direct effect of salmon fishing in the river, but with a twist. "We found that a special net type, a salmon weir, that accounts for majority of net catches, is capturing predominantly smaller fish, although net fishing is often assumed to catch larger fish," says Jaakko Erkinaro, a Research Professor at Natural Resources Institute Finland (Luke). Indigenous knowledge provided an answer for this surprising result: "We discussed with local Sámi fishers who have fished with salmon weir for decades, and they explained that, compared to other net types, salmon weir has smaller mesh size, is used later in the season and mostly in shallower waters, which increases the catch of smaller salmon. This likely explains the result," Erkinaro continues. Weir fishing has decreased in recent years, and simultaneously, the proportion of small, early maturing fish in spawning populations has increased.

"Our finding that there are different forms of fishing acting in opposite directions at different phases of their life-cycle highlights new challenges for salmon management, but also the value of having unique long-term data series at our disposal," noted Erkinaro.

The research team examined scale samples from salmon over a 40-year period and linked the variation of a gene that determines salmon reproduction age and size, with the effects of different fishing methods.

The scale samples used for the study came from a unique long-term scale archive maintained by Luke. The archive keeps samples from more than 150,000 salmon individuals collected by trained, volunteer fishers since the 1970's from River Teno, one of the most prolific salmon rivers in Europe. The scales were used to determine the age structure of the [salmon](#) population. They were also the source of DNA for genetic analysis.

**More information:** Y. Czorlich et al, Rapid evolution in salmon life-history induced by direct and indirect effects of fishing, *Science* (2022). [DOI: 10.1126/science.abg5980](https://doi.org/10.1126/science.abg5980).  
[www.science.org/doi/10.1126/science.abg5980](https://www.science.org/doi/10.1126/science.abg5980)

Provided by Natural Resources Institute Finland

Citation: Wild salmon are getting smaller and it might be linked to aquaculture (2022, February 25) retrieved 16 May 2024 from <https://phys.org/news/2022-02-wild-salmon-smaller-linked-aquaculture.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--