

# **Advances in micro-computed tomography**

#### March 8 2022



Julia Herzen, Professor of Biomedical Imaging Physics at TUM, working together with her team at the micro-computed tomography scanner. Credit: René Lahn

Researchers in biomedical physics and biology have significantly improved micro-computed tomography, more specifically imaging with phase contrast and high brilliance X-ray radiation. They have developed a new microstructured optical grating and combined it with new analytical algorithms. The new approach makes it possible to depict and analyze the microstructures of samples in greater detail, and to investigate a particularly broad spectrum of samples.

Micro-computed tomography (micro-CT) is an imaging method which



generates detailed three-dimensional <u>images</u> of the internal structure of samples with small dimensions. Researchers in biology, medicine or material sciences can use this method to obtain information on the structure and characteristics of tissue and material samples which are important in diagnoses and other analyses.

Micro-CT is based on X-ray images which are reconstructed to form a three-dimensional image. Depending on the sample, different X-ray imaging methods are used in order to achieve the most accurate depiction possible. Here the key parameters are resolution, <u>contrast</u> and the sensitivity of the method used.

# X-ray imaging with phase contrast

X-ray imaging with <u>phase contrast</u> is particularly well-suited for investigating soft tissue. The method employs the refraction of the Xrays caused by the sample's structures in order to obtain contrast for these structures and thus to depict soft tissue in greater detail than it is possible with conventional X-ray methods.

In many phase-contrast methods, optical components modulate the Xrays on their way to the detector, resulting in what is referred to as a diffraction pattern at the detector. "When comparing this pattern with and without the sample in the X-ray beam, the refraction of the X-rays on the sample provides information about its characteristics," says Julia Herzen, professor of Biomedical Imaging Physics at the Technical University of Munich (TUM).

Until now inefficient structures such as sandpaper and absorption masks have been used for this type of modulation, but in the meantime a variety of optical gratings are available. "The function of the new optical gratings resembles that of small lenses. The gratings focus the X-rays to form tiny points. This renders the differences in intensity with and



without the sample much clearer and makes it possible to visualize even minute differences in the tissue in greater detail," says Prof. Herzen.

## High contrast, high resolution and high sensitivity

Physicist Julia Herzen and her team have now introduced a new method for micro-CT with phase contrast using high-brilliance X-ray radiation. The technology is based on a newly developed optical grating referred to as a Talbot Array Illuminator. This new optical element is comparatively easy to produce, is resilient to X-ray radiation and can be used with different energies. This establishes the technically necessary prerequisites for high contrast. The new method enables more efficient use of the radiation dose than with ordinary modulators such as sandpaper and significantly reduces scan times.

"By combining our newly developed Talbot Array Illuminator with new analysis software optimized for the purpose, we've been able to significantly improve imaging and analysis with micro-CT. The new technology is more sensitive than comparable methods in this field. At very high resolutions, it allows to depict soft tissue with higher contrast than previously. High sensitivity is particularly important for example in order to detect fine differences within soft tissue," says Prof. Herzen.

#### Analysis of a broad spectrum of samples

The new technology can be used to investigate a particularly broad spectrum of samples. Researchers can even simultaneously depict materials of greatly differing compositions, for example water and oil embedded in stone, which was not possible in the past using conventional methods. This provides crucial advantages over conventional methods not only in medicine and biology, but also opens up new application possibilities in material sciences, for example in geology.



### Quantitative analysis is possible

"In contrast to previous approaches, our new method also makes quantitative analysis possible. We can make and compare absolute measurements of the electron density of samples, without the need for any assumptions about the samples," explains Prof. Herzen. Further studies will investigate the potential of this new option in a variety of applications.

The research was published in Optica.

**More information:** Alex Gustschin et al, High-resolution and sensitivity bi-directional x-ray phase contrast imaging using 2D Talbot array illuminators, *Optica* (2021). <u>DOI: 10.1364/OPTICA.441004</u>

Provided by Technical University Munich

Citation: Advances in micro-computed tomography (2022, March 8) retrieved 24 May 2024 from <u>https://phys.org/news/2022-03-advances-micro-computed-tomography.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.