

## **Corals can be 'trained' to tolerate heat stress, study finds**

March 3 2022



A subset of corals showing healthy coloration at the beginning of the heat-stress assay following either the laboratory control or variable temperature treatment. Photos such as this were taken daily in order to track the progression of bleaching during the heat-stress assay. Credit: Amanda Kirkland

## A new study led by researchers at the University of Miami (UM)



Rosenstiel School of Marine and Atmospheric Science found that corals that underwent a stressful temperature treatment in the laboratory for 90 days were more tolerant to increased water temperatures.

These findings offer coral restoration scientists with a new approach to potentially increase the success rate of planting nursery-raised staghorn coral onto degraded reefs as climate change continues to <u>warm ocean</u> <u>temperatures</u>, resulting in more frequent coral bleaching events. Staghorn coral (Acropora cervicornis) has died off throughout South Florida and the Caribbean, and is listed as "threatened" on the Endangered Species Act.

While previous "stress-hardening" experiments on corals have utilized exposures to short-term temperatures, the UM Rosenstiel School team assessed the effect of a long-term, variable <u>treatment</u> where temperatures reached a stressful level for a brief period of time, twice per day.

"This 'training' regime is akin to an athlete preparing for a race," said the study's lead author Allyson DeMerlis, a Ph.D. student at the UM Rosenstiel School. "We were able to demonstrate that this temperature treatment can boost the corals' stamina to heat stress."

To conduct the experiment, DeMerlis and scientists at the National Oceanic and Atmospheric Administration's (NOAA) Atlantic Oceanographic and Meteorological Laboratory, and UM's Cooperative Institute for Marine and Atmospheric Studies, collected coral fragments from six distinct genetic individuals of Caribbean staghorn coral from the UM Rosenstiel School's Rescue a Reef coral nursery and randomly assigned them to one of three groups: (1) field control, (2) <u>laboratory</u> control, and (3) variable temperature treatment. The laboratory control and variable temperature-treated corals were subjected to a three-month treatment period where the laboratory controls were kept at a constant



28 degrees Celsius while the variable temperature regime corals were subjected to fluctuating temperatures between 28 to 31 degrees Celsius, twice daily for three months.

The scientists then measured bleaching progression photographically as well as the number of days that a coral endured thermal stress before bleaching. They found that the variable temperature treatment significantly improved coral endurance in thermal stress, on the order of several days, in comparison to the untreated corals. In addition, they found that untreated corals were more likely to quickly succumb to disease-like signs of tissue loss.



Staghorn coral (Acropora cervicornis) maintained in an aquarium in the Experimental Reef Laboratory. Credit: Allyson DeMerlis,



The findings demonstrate the benefit of using a variable temperature treatment in the laboratory setting for maintaining staghorn coral over the traditional static temperatures. This may be translated in the field for restoration practitioners, specifically for identifying locations where their coral nurseries and outplanting sites can be exposed to more fluctuating temperatures.

"We have unfortunately reached the point where active intervention and restoration are necessary to ensure that valuable coral reefs are able to persist for generations to come," said Ian Enochs, senior author of the study and a <u>coral</u> scientist at NOAA's Atlantic Oceanographic and Meteorological Laboratory, Ocean Chemistry and Ecosystems Division. "We want to increase the efficiency and efficacy of these efforts, and ultimately ensure that the corals that are placed back out on a reef have the greatest chance of enduring the stressful conditions they will face in the future."

"Our findings bring a glimmer of hope to the uncertain future of corals, as we identified a treatment in which we can enhance their tolerance to heat stress," said DeMerlis.

The study, titled "Pre-exposure to a variable <u>temperature</u> treatment improves the response of Acropora cervicornis to acute thermal stress," was published Feb. 23 in the journal *Coral Reefs*.

**More information:** Allyson DeMerlis et al, Pre-exposure to a variable temperature treatment improves the response of Acropora cervicornis to acute thermal stress, *Coral Reefs* (2022). DOI: 10.1007/s00338-022-02232-z

Provided by University of Miami



Citation: Corals can be 'trained' to tolerate heat stress, study finds (2022, March 3) retrieved 3 May 2024 from <u>https://phys.org/news/2022-03-corals-tolerate-stress.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.