


Controlling how fast graphene cools down

March 4 2022



Graphene charge carriers lying on different energetic levels represented by the

Dirac cones, which, depending on the number of charge carriers, are occupied up to the neutrality point (blue level on the left cone) or well into the conduction band (blue level on the right cone). In the two cases, the photoexcited charge carriers relax with faster (left side) or slower (right side) dynamics. Credit: Politecnico di Milano - CNR

Graphene is the thinnest material ever produced, with the thickness of a single atomic layer. Thinner than a billionth of a meter, it is able to efficiently absorb light from the visible to the infrared through the photoexcitation of its charge carriers. After light absorption, its photoexcited charge carriers cool down to the initial equilibrium state in a few picoseconds, corresponding to a millionth of a millionth of a second. The remarkable speed of this relaxation process makes graphene particularly promising for a number of technological applications, including light detectors, sources and modulators.

A recent study published in *ACS Nano* has shown that the [relaxation](#) time of graphene charge carriers can be significantly modified by applying an external electrical field. The research was conceived within an [international collaboration](#) between the CNR-IFN, Politecnico di Milano, the University of Pisa, the Graphene Center of Cambridge (UK) and ICN2 of Barcelona (Spain).

"The change in the relaxation time of charge carriers in graphene that we have observed, demonstrates an unprecedented level of control on the optical response of a crystal and allows to obtain a large variety of behaviors using a single material" says Eva Pogna, researched from CNR-IFN, first author of the work.

This work paves the way to the development of devices that exploit the control of the relaxation time of charge carriers to support novel

functionalities. For example, if graphene is used as saturable absorber in a laser cavity to generate ultrashort light pulses, by changing the relaxation time of the charge carriers, we can control the duration of the output pulses.

"The specific device that we have used to study graphene, proved to be crucial to observe the strong tunability of its optical properties with the external electric field, allowing to change the number of charge carriers over a broad range by exploiting ionic liquid gating, which is a state-of-the-art technology introduced to study superconductors" explains Andrea Ferrari, director of the Graphene Center in Cambridge.

The graphene-based device has been studied by ultrafast spectroscopy, which allowed to monitor the variation of the relaxation time of the charge carriers.

"This work represents the latest step of a long-standing research collaboration devoted to the study of the ultrafast [carrier](#) dynamics in [graphene](#), aimed at exploring the great potential of this fascinating material" as added by Klaas-Jan Tielrooij, leader of the Ultrafast Dynamics in Nanoscale Systems group at ICN2.

"This discovery is of large interest for a number of technological applications, ranging from photonics, for pulsed laser sources or optical limiters that prevent optical components damaging, to telecommunication, for ultrafast detectors and modulators" concludes Giulio Cerullo, professor of the Physics Department of Politecnico di Milano.

More information: Eva A. A. Pogna et al, Electrically Tunable Nonequilibrium Optical Response of Graphene, *ACS Nano* (2022). [DOI: 10.1021/acsnano.1c04937](https://doi.org/10.1021/acsnano.1c04937)

Provided by Politecnico di Milano

Citation: Controlling how fast graphene cools down (2022, March 4) retrieved 2 May 2024 from <https://phys.org/news/2022-03-fast-graphene-cools.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.