


Hundreds of gene regulatory motifs cooperate and conflict to make brain cells

March 25 2022



Experimental design. a Computational framework to select regions and perturbation sites. b Heatmap of motif instances in the assayed regions (left); distribution of the number of motifs perturbed in each region (top right); distribution of the number of regions harboring each motif (bottom right). c Library design; selected regions were included in their wild-type (WT) form, selected motifs were perturbed (by altering the sequence in the predicted motif site) using three perturbation methods individually (PERT single) as well as in combination with other perturbations in selected cases (PERT double). Random sites were perturbed (RAND) and the entire WT sequence was scrambled as negative controls (SCRAM) for each WT sequence. d The designed sequences

were synthesized and cloned into the lentiMPRA vector and associated with 15-bp barcodes. ARE antirepressor element, BC barcode. Reporter, EGFP enhanced green fluorescent protein, LTR long terminal repeat, mP minimal promoter, WPRE Woodchuck Hepatitis Virus Posttranscriptional Regulatory Element. e lentiMPRA libraries were infected into hESCs and following 3 days, we induced neural differentiation via dual-SMAD inhibition and obtained DNA and RNA at seven time points (0, 3, 6, 12, 24, 48, and 72 h). f Association between barcodes and designed sequences, and the number of barcodes observed in DNA and RNA sequencing was determined using MPRAflow. Differential analysis between WT and PERT activity to determine motif regulatory effect over time was assessed using MPRAnalyze. Source data are provided as a Source Data file. Credit: *Nature Communications* (2022). DOI: 10.1038/s41467-022-28659-0

DNA is considered the blueprint of our human bodies. It codes for the tens of thousands of genes that determine whether a cell transforms into a brain cell, heart cell or even cancer cell. Other DNA sequences, known as enhancers, determine if genes are activated or inactivated. In a new study seen in *Nature Communications*, ASHBi Associate Professor Fumitaka Inoue and colleagues in the United States report the "massively parallel reporter perturbation assay" to study how enhancers are fine-tuned over time during the generation of brain cells.

Enhancers determine when, where and to what degree a gene is turned on. Enhancers themselves are regulated by [transcription factors](#), which target regulatory motifs in the enhancer sequence. Thus, interactions between enhancers and transcription factors are critical for enhancing or suppressing the genes that determine a cell's fate and behavior.

"Enhancers have important effects on [gene expression](#) and thus determine whether cells will grow into different organs. Despite their consequences, however, we understand relatively little about how

enhancers function," explains Inoue.

A number of biochemical assays are available to study enhancers, but massively parallel reporter assays (MPRA) allow for the [quantitative analysis](#) of thousands of enhancers at one time.

"MPRA are useful for studying the interplay between [enhancers](#) and transcription factors. To extract even more information from the assays, we modified the assays to introduce perturbations in the regulatory motifs where transcription factors bind," continues Inoue.

These perturbations were introduced in multiple regulatory motifs in thousands of enhancer sequences. MPRA was then applied to study the effects of the perturbations on the enhancer activity and thus systematically identify which regulatory motifs are important for gene activation or inactivation.

To validate their massively parallel reporter perturbation assay, the team applied it to the differentiation of human embryonic stem cells into neural cells. They found 598 regulatory motifs that either enhanced or dampened transcription.

"We broke these motifs into four groups. Essential and contributing motifs either were required or contributed to the activation. Likewise, silencing and inhibiting motifs were required or contributed to dampening transcription," says Inoue.

Further investigation revealed that among the [genes](#) enhanced or dampened, many are involved in promoting neural differentiation or stem cell pluripotency. Moreover, the assay clarified that in most cases the regulatory motifs determined whether transcription was enhanced or dampened, but the timing of the perturbation determined the magnitude of the effect. Overall, says Inoue, the findings demonstrate an intricate

network of regulatory sequences that interact with one another to achieve neural differentiation.

"The biggest accomplishment of the assay is the number of sequences we could analyze and perturbations we could apply. As a result, we demonstrated the complexity of many models for how transcription factors regulate gene expressions," he said.

More information: Anat Kreimer et al, Massively parallel reporter perturbation assays uncover temporal regulatory architecture during neural differentiation, *Nature Communications* (2022). [DOI: 10.1038/s41467-022-28659-0](https://doi.org/10.1038/s41467-022-28659-0)

Provided by Kyoto University

Citation: Hundreds of gene regulatory motifs cooperate and conflict to make brain cells (2022, March 25) retrieved 23 April 2024 from <https://phys.org/news/2022-03-hundreds-gene-regulatory-motifs-cooperate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.