

# A not-so-solid lake ice season

March 1 2022

---



Lake ice cover in Green Lakes Valley, Colorado, has decreased by 24 days since 1983. Credit: Kelly Loria CC BY-SA 4.0.

Mountain lakes provide freshwater for roughly half the world's population. Yet a shifting climate could impact the reliability of this

resource by altering the temperature patterns that govern lake conditions and influence lake biota.

Lake ice phenology, or the timing of lake ice cover, tell researchers how a lake is responding to climate shifts. But long-term data sets are necessary to track slow, imperceptible changes in alpine systems, scientists say.

A study by researchers affiliated with the U.S. National Science Foundation Niwot Ridge Long-Term Ecological Research site reported a 24-day decrease in lake ice cover duration in Green Lakes Valley, Colorado, since 1983. Earlier spring thaw dates (nine days earlier) and especially later autumn freeze dates (14 days later) contributed to the decrease in ice cover.

That translates to 20% more open water time and represents an annual rate of ice cover loss that is approximately 50% faster than the Northern Hemisphere average. The results are published in *Limnology and Oceanography Letters*.

"One of our ideas is the 'extended summer,'" says Kyle Christianson of the University of Colorado Boulder and lead author of the study.

"There's an earlier start to melting in the spring and later start to ice in the fall. A small increase in the extension of the open water season increases primary productivity, with far-reaching ecological implications."

Alpine lakes are not isolated systems; changes in these lakes likely indicate watershed-level changes, too. "The extended summer has led to terrestrial vegetation increases and some vertical movement of tree species, contributing to watershed-level shifts," says Christianson.

Such drastic seasonal shifts have social and ecological implications.

Warmer, longer summers support more biological productivity, which requires water treatment facilities to re-think their purification processes.

Lake ice has been studied in Green Lakes Valley at the Niwot Ridge LTER site for 51 years, the longest-running high elevation dataset on lake ice globally. Christianson and collaborators studied five of the seven Green Lakes Valley lakes. They focused on a 36-year window in the data where all five lakes had continuous recordings of ice clearance and formation.

**More information:** Kyle R. Christianson et al, On thin ice: Linking elevation and long-term losses of lake ice cover, *Limnology and Oceanography Letters* (2021). [DOI: 10.1002/lol2.10181](https://doi.org/10.1002/lol2.10181)

Provided by National Science Foundation

Citation: A not-so-solid lake ice season (2022, March 1) retrieved 18 August 2024 from <https://phys.org/news/2022-03-not-so-solid-lake-ice-season.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--