


## Study reveals lunar regolith evolution process on Chang'e-4 landing area

## March 31 2022, by Li Yuan



Evolution of lunar regolith in the Chang'e-4 landing area. (a), (b) and (c) are the spatial distribution of lunar regolith thickness at 3.0 Ga, 2.0 Ga and 1.0 Ga, respectively; (d), (e) and (f) are the profiles of lunar regolith thickness at the



corresponding time of horizontal 200 m; (g), (h) and (i) are the profiles of lunar regolith size at the corresponding time of horizontal 200 m. The units are all in meters. Credit: AIR

A research team led by Prof. Di Kaichang from the Aerospace Information Research Institute (AIR) of the Chinese Academy of Sciences (CAS) and their collaborators have established an evolution model with simple craters in the Chang'e-4 landing area. The research results reveal the evolution process of lunar regolith and quantify the overturning and mixing effect of small simple impact events on lunar regolith.

The study was published in *Icarus*.

China's Chang'e-4 probe, launched on Dec. 8, 2018, made the first-ever soft landing on the Von Karman Crater in the South Pole-Aitken Basin on the far side of the moon on Jan. 3, 2019.

Lunar regolith, a layer of unconsolidated debris covered on the surface of the moon, is generally supposed to be the main product of impacts from meteorites or micro meteorites and high-energy particle bombardment. Lunar regolith evolves under the effect of impact gardening and the process is significant to the research of lunar stratigraphy, lunar impact history and the energy distribution on the lunar surface.

The research team interpreted the <u>impact craters</u> from high-resolution remote sensing images, and obtained a group of random impact events in the landing area according to the Neukum production function and chronology function of lunar crater.



Based on the established impact events, the team analyzed the evolution process of lunar surface materials under the effect of impact gardening.

They first analyzed the impact excavation of a single impact event and calculated the superposition of multiple impact events. Then, the impact events were generated one-by-one according to the production function of the lunar impact crater, and the calculation was carried out step-by-step.

Eventually, the spatial distribution of the lunar regolith thickness and the fragmental size in the Chang'e-4 landing area at different ages were obtained.

The study indicates that the formation of lunar regolith is controlled by larger impact events, while the surficial overturning and mixing of lunar regolith are mainly caused by smaller impact events.

In addition, it also reveals that the lunar regolith is mostly formed in early due to the high impact flux at that time. For example, the absolute model age of the Chang'e-4 landing area is about 3.5 Ga, while the lunar regolith had mostly been formed before 3.0 Ga and changed slowly afterwards.

Based on in situ data from Chang'e-4 probe, scientists can determine the thickness of the <u>regolith</u> and reveal the fine subsurface structures and evolutionary history of the probe's landing area.

With the model, scientists speculated a possible scenario of the regional <u>lunar regolith</u> formation and evolution process, which will provide reference for future missions of lunar surface exploration and sampling.

**More information:** Ke Shi et al, The gardening process of lunar regolith by small impact craters: A case study in Chang'E-4 landing area,



Icarus (2022). DOI: 10.1016/j.icarus.2022.114908

## Provided by Chinese Academy of Sciences

Citation: Study reveals lunar regolith evolution process on Chang'e-4 landing area (2022, March 31) retrieved 21 May 2024 from https://phys.org/news/2022-03-reveals-lunar-regolith-evolution-change-.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.