

After 20 years of studying how ecosystems absorb carbon, why we're worried about a tipping point of collapse

March 25 2022, by Caitlin Moore, David Campbell, Helen Cleugh, Jamie Cleverly, Jason Beringer, Lindsay Hutley and Mark Grant



Each hectare of Australia's savanna's sequesters, on average, 3.4 tonnes of carbon every year. Credit: Bryn Pinzgauer/Wikimedia, <u>CC BY-SA</u>



From rainforests to savannas, ecosystems on land absorb <u>almost 30%</u> of the carbon dioxide human activities release into the atmosphere. These ecosystems are critical to stop the planet warming beyond 1.5°C this century—but climate change may be weakening their capacity to offset global emissions.

This is a key issue that OzFlux, a research network from Australia and Aotearoa New Zealand, has been investigating for the past 20 years. Over this time, we've identified which <u>ecosystems</u> absorb the most carbon, and have been learning how they respond to <u>extreme weather</u> and <u>climate events</u> such as drought, floods and bushfires.

The biggest absorbers of atmospheric <u>carbon dioxide</u> in Australia are savannas and temperate forests. But as the effects of <u>climate change</u> intensify, ecosystems such as these are at risk of reaching tipping points of collapse.

In our latest <u>research paper</u>, we look back at the two decades of OzFlux's findings. So far, the ecosystems we studied are showing resilience by rapidly pivoting back to being <u>carbon sinks</u> after a disturbance. This can be seen, for example, in leaves growing back on trees soon after bushfire.

But how long will this resilience remain? As climate change pressures intensify, evidence suggests carbon sinks may lose their ability to bounce back from climate-related disasters. This reveals vital gaps in our knowledge.

Australian ecosystems absorb 150 million tons of carbon each year

Between 2011 and 2020, land-based ecosystems sequestered 11.2 billion



tons (29%) of global CO₂ emissions. To put this into perspective, that's roughly similar to the amount China emitted in 2021.

OzFlux has enabled the first comprehensive assessment of <u>Australia's</u> <u>carbon budget</u> from 1990 to 2011. This found Australia's land-based ecosystems accumulate some 150 million tons of CO_2 each year on average—helping to offset national fossil fuel emissions by around one third.

For example, every hectare of Australia's <u>temperate forests</u> absorbs 3.9 tons of carbon in a year, <u>according to OzFlux data</u>. Likewise, every hectare of Australia's savanna absorbs 3.4 tons of carbon. This is about 100 times larger than a hectare of Mediterranean woodland or shrubland.

But it's important to note that the amount of carbon Australian ecosystems can sequester fluctuates widely from one year to the next. This is due to, for instance, the <u>natural climate variability</u> (such as in La Niña or El Niño years), and disturbances (such as fire and land use changes).

In any case, it's clear these ecosystems will play an important role in Australia reaching its target of net-zero emissions by 2050. But how effective will they continue to be as the climate changes?

How climate change weakens these carbon sinks

Extreme climate variability—flooding rains, droughts and heatwaves—along with bushfires and <u>land clearing</u>, can weaken these carbon sinks.



Tonnes of carbon absorbed by one hectare of Australian ecosystems, every year

Based on two decades of OzFlux's findings



Credit: Beringer et al, 2016

While many Australian ecosystems show resilience to these stresses, we found their <u>recovery time</u> may be shortening due to more frequent and <u>extreme events</u>, potentially compromising their long-term contribution towards offsetting emissions.

Take bushfire as an example. When it burns a forest, the carbon stored



in the plants is released back into the atmosphere as smoke—so the ecosystem becomes a carbon source. Likewise, under drought or heatwave conditions, water available to the roots becomes depleted and limits photosynthesis, which can tip a forest's carbon budget from being a sink to a carbon source.

If that drought or heatwave endures for a long time, or a bushfire returns before the forest has recovered, its ability to regain its carbon sink status is at risk.

Learning how carbon sinks may shift in Australia and New Zealand can have a global impact. Both countries are home to a broad range of climates—from the wet tropics, to the Mediterranean climate of southwest Australia, to the temperate climate in the southeast.

Our unique ecosystems have evolved to suit these diverse climates, which are underrepresented in the global network.

This means long-term ecosystem observatories—<u>OzFlux</u>, along with the <u>Terrestrial Ecosystem Research Network</u>—provide a vital natural laboratory for understanding ecosystems in this era of accelerating climate change.

Over its 20 years, OzFlux has made crucial contributions to the international understanding of climate change. A few of its major findings include:

- The 2011 La Niña event led to a greening of interior Australia, with ecosystems flourishing from increased water availability.
- Heatwaves can negate the carbon sink strength of our ecosystems, and even lead to carbon emissions from plants .
- <u>Land clearing</u> and the <u>draining of peatland</u> systems add to Australia's and New Zealand's greenhouse gas emissions.



Critical questions remain

Plans in Australia and New Zealand to reach net zero emissions by 2050 strongly depend on the ongoing ability for ecosystems to sequester emissions from industry, agriculture, transport and the electricity sectors.

While some management and technological innovations are underway to address this, such as in the agricultural sector, we need long-term measurements of carbon cycling to truly understand the limits of ecosystems and their risk of collapse.

Indeed, we're already in uncharted territory under climate change. Weather extremes from heatwaves to <u>heavy rainfall</u> are becoming more frequent and intense. And CO_2 levels are more than <u>50% higher</u> than they were 200 years ago.

So while our ecosystems have remained a net sink over the <u>last 20 years</u>, it's worth asking:

- Will they continue to do the heavy-lifting required to keep both countries on track to meet their climate targets?
- How do we protect, restore and sustain the most vital, yet vulnerable, ecosystems, such as "<u>coastal blue carbon</u>" (including seagrasses and mangroves)? These are critical to nature-based solutions to climate change.
- How do we monitor and verify national carbon accounting schemes, such as Australia's <u>Emissions Reduction Fund</u>?

Critical questions remain about how well Australia's and New Zealand's ecosystems can continue storing CO₂.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.



Provided by The Conversation

Citation: After 20 years of studying how ecosystems absorb carbon, why we're worried about a tipping point of collapse (2022, March 25) retrieved 25 June 2024 from https://phys.org/news/2022-03-years-ecosystems-absorb-carbon-collapse.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.