

Adriatic ecosystems withstand major climate shifts but wither under human impact

April 18 2022, by Jerald Pinson



Adriatic mollusks have proven to be incredibly resilient to climate change, weathering the last ice age with nearly no long-term effects. But human activity over the last few centuries threatens to destabilize mollusk communities in the region, with negative impacts for the marine ecosystems they occupy. Credit: Fabio Negri



An analysis of more than 70,000 fossils indicates that mollusk communities were incredibly resilient to major climatic shifts during the last ice age.

Scientists from the Florida Museum of Natural History and several European research institutions tracked the history of Adriatic <u>ecosystems</u> through two warm periods that bookend the most recent glacial expansion. Their results show that major changes in temperature, salinity and sea level had much less of an impact on <u>mollusk</u> communities than the current environmental crisis caused by <u>human activity</u> in the region.

"It is sobering to consider that about 120,000 years of major <u>climate</u> <u>change</u> did not affect these ecosystems nearly as much as the humaninduced changes of the last few centuries," said senior author Michał Kowalewski, the Florida Museum Thompson chair of Invertebrate Paleontology.

Researchers have known for some time that modern Adriatic ecosystems are considerably altered compared to historical baselines.

"There are multiple human-driven stressors on these ecosystems, such as changes in land use that increase sedimentation rates," said co-author Rafał Nawrot, a postdoctoral researcher at the University of Vienna, formerly with the Florida Museum. "This has occurred as far back as the Roman Empire, when increased agriculture led to higher rates of erosion."

While prior civilizations along the Italian peninsula have left a notable signature on Adriatic ecosystems, Nawrot explains that most of the changes have occurred within the last century. The influx of fertilizers into rivers and estuaries have sparked runaway reactions that deplete oxygen in marine and freshwater environments. Pollution from towns and cities creates a toxic mélange for <u>sea life</u>, and cargo vessels litter



international shipping lanes with invasive stowaway species that disrupt native ecosystems. Perhaps most detrimental to Adriatic mollusks, <u>commercial fisheries</u> drag nets across the sea floor, scouring the basin for bottom-dwelling fish and bivalves.



The researchers obtained fossils from sediment cores drilled in terrestrial and deep-sea environments. Credit: Fabiano Gamberi

"The Adriatic Sea is the most heavily trawled area in the world," Nawrot said, citing a <u>broad study</u> published earlier this year.

What scientists have lacked, up until now, was data on natural disturbances in the Adriatic's recent past they could use to assess the significance of current ecological changes.



"By looking at the <u>fossil record</u>, you can reconstruct a range of natural variability. If the present-day community falls outside that range, it's probably because of us," said lead author Daniele Scarponi, an associate professor at the University of Bologna.

To fill the gap in the region's history, the researchers sieved long, vertical sediment cores for marine mollusk fossils dating back to the Late Pleistocene.

The ice ages were a turbulent time for global marine ecosystems. Sea levels rose and fell as water from the world's oceans was cyclically released from and trapped in massive continental glaciers. All told, there were about 17 ice ages during the Pleistocene, starting approximately 2.5 million years ago and ending with the final retreat of the glaciers just 12,000 years before present.

The changes wrought by the onset of the last ice age are especially evident in the Adriatic Sea, which is mostly shallow across its northern extent. When sea levels dropped by about 400 feet at the peak of the ice age, the northern Adriatic nearly ceased to exist, its shores retreating over 150 miles south toward the Mediterranean.





Map: The northern Adriatic is mostly shallow, with depths of less than 200 feet. As sea levels rose and fell during the glaciations of the Pleistocene, large swaths of land were inundated and exposed. Credit: Alfonso Di Pasquale (1961) / The European Commission, CC-BY-4.0. Globe: Www.Onestopmap.Com / CC-BY

"It would have been possible to walk from modern-day Italy to Croatia," Scarponi said. "The entire northern part of the basin was exposed and transformed into an immense lowland plain."

By studying fossils preserved before, during and after the last ice age, the researchers could directly observe the changes to mollusk communities. Their results show that as oceans receded and cooled, mollusks that are today restricted to colder regions flourished in the Adriatic while warmth-loving species declined.



The researchers note, however, that despite the strong shift in numbers, very few species disappeared altogether. "The main difference between mollusk communities through the glacial/interglacial cycles isn't extinction or the appearance of new species, but rather changes in relative abundance," Nawrot said.

When the glaciers receded and the Adriatic swelled to its modern boundaries, mollusk species reverted back to their previous abundance patterns. If scientists hadn't known there was an ice age in between, it would have seemed like nothing much had happened at all based solely on mollusks.

"Our fossil analyses show that communities of mollusk species along northern Adriatic coasts essentially re-assembled into a nearly identical image of themselves when the sea returned," Kowalewski said.

By itself, the finding that mollusks are resilient to climate change is good news. Modern marine communities are already contending with increasing ocean temperatures associated with global warming caused by humans.





Mollusk species underwent a stark change in abundance during the most recent ice age, but a nearly identical species assemblage returned as temperatures warmed and sea levels rose. Credit: Daniele Scarponi

"Temperatures during the last interglacial were actually a few degrees



higher than they are at present, and yet we see the same associations of mollusks," Scarponi said. "That means near-shore mollusk communities will likely be resilient to a slight increase in temperature going forward."

But a warmer world brings with it a unique set of compounding problems, warns Nawrot. "Many stressors, like anoxia and the effects of invasive species will only intensify with warming, even if higher temperatures alone wouldn't be a huge deal," he said.

The future of marine ecosystems in the Adriatic and throughout the world's oceans remains an open-ended question, one the study authors maintain will require a variety of strategies to solve. "We need international policies addressing global climate change, but studies like these show we also need actions that mitigate local and regional threats," Nawrot said.

More information: Daniele Scarponi et al, Resilient biotic response to long-term climate change in the Adriatic Sea, *Global Change Biology* (2022). DOI: 10.1111/gcb.16168

Provided by Florida Museum of Natural History

Citation: Adriatic ecosystems withstand major climate shifts but wither under human impact (2022, April 18) retrieved 2 May 2024 from <u>https://phys.org/news/2022-04-adriatic-ecosystems-major-climate-shifts.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.