

Eelgrass beds important for resilience of fish stocks

April 20 2022, by Sissel Eikeland



The common eelgrass or seawrack—Zostera subg. Zostera marina Linnaeus, 1753. Photograph taken in Puget Sound, Edmonds, Washington state, USA (June 7, 2020). Credit: John Brew/Wikimedia Commons, <u>CC BY 4.0</u>



A new study from UiA shows that a network of eelgrass beds is important for maintaining the genetic diversity of fish stocks, and thereby making them more resilient against changes in the environment, like climate change.

"The study shows that eelgrass is important for the different species. If the distance between eelgrass areas becomes too large for organisms to cross, local <u>fish stocks</u> will not exchange individuals or genes. The stocks thus become less resilient to overfishing and <u>environmental changes</u>," says Halvor Knutsen.

He is director of the Center for Coastal Research (CCR) at UiA and works at UiA and the Institute of Marine Research in Flødevigen in Arendal. Knutsen carried out the study in collaboration with marine scientists from CCR and the Institute of Marine Research. The study was recently published in the international journal *Molecular Ecology*.

From the Swedish border to Ålesund in western Norway, coastal fish species such as broadnosed pipefish, corkwring wrasse and black goby have been compared. The aim was to find common features in how the species experience barriers and how they are prevented from mixing with other populations.

Eelgrass grows in patches along the entire Norwegian coast and form habitats for large ecosystems in the sea. A number of fish species live in <u>eelgrass beds</u>. When eelgrass areas are close to each other, the fish stocks can more easily mix. Eelgrass beds are important for fish populations to be able to mix.

"Eelgrass beds tie the populations together. We know that areas with good genetic variation provide more resilient stocks," Knutsen says.

Loss of contact with each other



The researchers are looking at an area on Jæren, for example, where natural sandbanks stretch over several kilometers. There, the constant waves make it difficult for eelgrass to grow.

"Large distances and the absence of eelgrass create longer distances between the populations and can lead to them losing contact with each other," Knutsen says.

In many areas, the researchers also see that eelgrass beds have to give way to jetties, artificial beaches, or other interventions along the coast. This affects the development of the species in the sea.

"Where they don't have contact with each other, they are more exposed to loss of genetic variation. This weakens the fish species and makes it less resistant to environmental changes," Knutsen says.

Genetic mapping

The study is part of Ecogenome, a research project funded by the Research Council of Norway. The research consists in the genetic mapping of a number of coastal species in Norway, including fish species, seaweed, and crustaceans. They also look at what steers the structuring into different stocks and the adaptations they make in the marine ecosystem.

"We map the genetic differences between the populations of several species. If we find a <u>coastal area</u> where several species experience a barrier, that may be a natural place to recommend a boundary for fish stock management," Knutsen says.

More information: Halvor Knutsen et al, Combining population genomics with demographic analyses highlights habitat patchiness and



larval dispersal as determinants of connectivity in coastal fish species, *Molecular Ecology* (2022). DOI: 10.1111/mec.16415

Provided by University of Agder

Citation: Eelgrass beds important for resilience of fish stocks (2022, April 20) retrieved 16 August 2024 from <u>https://phys.org/news/2022-04-eelgrass-beds-important-resilience-fish.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.