

Scientists crack egg forging evolutionary puzzle

April 11 2022



Cuckoo finch eggs laid by different females. Diversity of maternally inherited egg phenotypes within a single interbreeding species, the brood-parasitic cuckoo finch. Different cuckoo finch matrilineal mimics the eggs of different host species (here, tawny-flanked prinia and red-faced cisticola) and have further diversified to approximate the range of variable egg "signatures" within each host species, an anti-parasite adaptation that aids host parents in recognizing their own eggs. Credit: Claire N. Spottiswoode

As many humans prepare to unwrap their Easter eggs, scientists have solved one of nature's biggest criminal cases, an egg forgery scandal two million years in the making. Their findings suggest that the victims of this fraud may now be gaining the upper hand.

Around the world, many birds side-step the costs of parenthood by laying their eggs in the nest of other [species](#). This lifestyle, termed "[brood parasitism](#)", has many advantages but also presents challenges such as how to convince the other species to accept a foreign egg. Many [brood parasites](#) achieve this by mimicking the colors and patterns of their host's eggs, but some exploit the care of several different host species whose eggs all look different.

How then can a single brood-parasitic bird species simultaneously mimic the eggs of several different bird species to trick them into raising their young? And how do these parasitic forgers pass this ability on to their young despite interbreeding between birds raised by different hosts?

These questions have been puzzling scientists for more than a century. Now [genetic research](#) by an international team led by Professor Claire Spottiswoode from the University of Cambridge's Department of Zoology and the FitzPatrick Institute of African Ornithology, University of Cape Town; and Professor Michael Sorenson at Boston University, has made a major breakthrough, and their findings may be bad news for the egg forgers.

The study, published today in *PNAS*, focussed on the genetics of egg mimicry in the cuckoo finch, a species which adopts a brood-parasitic lifestyle and exploits many species of warbler across Africa. The research reveals that female cuckoo finches inherit their ability to mimic the appearance of their hosts' eggs from their mothers, via the female-specific W chromosome (analogous to the male-specific Y chromosome in humans).



Cuckoo finch and host chicks. Brood parasitism is costly for hosts because a cuckoo finch chick begs for food very vigorously as soon as it hatches, outcompeting the host parents' own chicks (here zitting cisticolas) which typically soon die of starvation. Credit: Claire N. Spottiswoode

Such '[maternal inheritance](#)' allows cuckoo finches to side-step the risk of inheriting the wrong mimicry genes from a father raised by a different host, and so has allowed distinct lineages of cuckoo finch females to evolve specialized egg mimicry of several different host species. Such mimicry dupes host parents into accepting a parasitic egg as their own rather than throwing it out of the nest, and so has been crucial to the success of these African birds.

But the researchers believe that this long-established 'genetic architecture' of maternal inheritance may come back to haunt the cuckoo finches. Dr. Spottiswoode said: "In this particular coevolutionary arms race between species, natural selection has created a double-edged sword.

"While maternal inheritance has allowed cuckoo finches to exploit multiple host species, it's likely to slow their ability to evolve counter-adaptations as their hosts evolve new defenses. In particular, parasites face a daunting challenge because some host species have in return evolved an astonishing diversity of egg color and pattern 'signatures', that help hosts to distinguish their own eggs from parasitic mimics."

The field data were collected at a study site in southern Zambia together with Dr. Wenfei Tong and Dr. Gabriel Jamie from the University of Cambridge and Ailsa Green, Silky Hamama, Ian Taylor and Collins Moya from the surrounding community in Zambia. Cuckoo finches in this area trick four different species of grass-warbler to devastating effect: if host parents fail to detect and remove a parasitic egg in their nest, the young cuckoo finch typically outcompetes the hosts' own hatchlings, which soon starve to death.

The team collected DNA samples from 196 cuckoo finches from 141 nests belonging to the four grass-warbler species and studied the majority by sequencing thousands of short segments across their

genomes.



A tawny-flanked prinia, a common host species of the cuckoo finch, captured in Zambia for genetic sampling with the help of field assistant Tom Hamusikili.
Credit: Claire N. Spottiswoode

In their fightback against the forgers, grass-warblers have become skilled quality controllers, rejecting eggs that differ from their own in color and pattern, and all four species have evolved the ability to deposit unique 'signatures' onto their own eggs to enhance their detection of intruders. Tawny-flanked prinias, for example, lay eggs with blue, white, red, or olive-green backgrounds overlaid with a variety of patterns.

Cuckoo finches have responded not only by evolving mimicry of the eggs of their several host species, but have also further diversified to mimic at least some of the signature-like variation seen in the eggs of different females within each host species. The team established that both abilities are handed down through maternal inheritance, finally validating a hypothesis first proposed in 1933 by ornithologists pondering how the common cuckoo in Europe was similarly able to mimic the eggs of several different host species.

Forgers facing an uncertain future?

The researchers believe that the cuckoo finches now face an uphill struggle because they cannot recombine the different forgery traits evolved by their separate family lines. For example, two different lineages of cuckoo finch mothers have evolved eggs with either blue or red backgrounds, as an evolutionary response to similar diversity in their tawny-flanked prinia hosts, but there is no evidence that they can create the precise mixture of pigments needed to produce the olive-green eggs that some host females can produce.

In a previous study, Professor Spottiswoode found that a growing proportion of eggs laid by tawny-flanked prinia hosts are olive-green, suggesting this is part of an accelerating evolutionary fightback. As expected, the team found that these host birds are passing down their anti-fraud 'egg signature' abilities through a different genetic process (bi-parental inheritance) to that used by the cuckoo finches.



Cuckoo finch egg in zitting cisticola nest. Cuckoo finch eggs closely mimic the colour and pattern of the eggs of each of their several host species, to trick host parents into accepting the parasitic egg as one of their own. Here a cuckoo finch has successfully had its egg (at left) accepted in the nest of a zitting cisticola (egg at right). Credit: Claire N. Spottiswoode

Spottiswoode said: "Cuckoo finches are missing out on a powerful source of evolutionary novelty and that could prove costly in this ongoing arms race. The way they inherit their ability to mimic host eggs has a downside by likely making the grass-warblers' defenses more effective, and constraining the parasite's ability to respond.

"We may see the emergence of unforgeable egg signatures which could force cuckoo finches to switch to other naïve [host species](#). Or the parasitic birds might become increasingly dependent on young host individuals that haven't yet learned their own signatures and are bad at spotting mismatched eggs."

The study argues that 'selection from host defenses drove [cuckoo](#) finches to transfer control of egg appearance to the maternally inherited part of the genome' at least 2 million years ago.

More information: Genetic architecture facilitates then constrains adaptation in a host–parasite coevolutionary arms race, *Proceedings of the National Academy of Sciences* (2022). [DOI: 10.1073/pnas.2121752119](#)

For more information, see africancuckoos.com

Provided by University of Cambridge

Citation: Scientists crack egg forging evolutionary puzzle (2022, April 11) retrieved 10 April 2024 from <https://phys.org/news/2022-04-scientists-egg-forging-evolutionary-puzzle.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
