


Hydrogen-tuned topological insulators may lead to new platforms in sustainable quantum electronics

May 4 2022



Rendering of hydrogenation resolving an outstanding challenge in chalcogenide classes of three-dimensional (3D) topological insulators and magnets. Credit: Lukas Zhao

A team led by City College of New York physicist Lia Krusin-Elbaum is behind research that could open a breadth of new quantum device platforms for harnessing emergent topological states for nano-spintronics and fault-tolerant quantum computing.

The group of physicists and chemists has invented a new facile and powerful technique that uses ionic hydrogen to reduce charge carrier density in the bulk of three-dimensional (3D) topological insulators and magnets. The result is that robust non-dissipative surface or edge quantum conduction channels can be accessed for manipulation and control. Their research, "Topological surface currents accessed through reversible hydrogenation of the three-dimensional bulk," appears in the journal *Nature Communications*.

The novel hydrogen-tuning technique of chalcogen-based topological materials and nanostructures implemented in a laboratory chamber uses insertion and extraction of ionic hydrogen from dilute aqueous hydrochloric acid (HCl) solution, which leaves the layered topological crystal structure as well as electronic bands intact and has an extra benefit of removing native surface oxide while passivating surfaces. In this process—which the City College team tests in the Krusin Lab for two-dimensional electrical transport—electrons are donated by a reversible binding of H^+ ions to chalcogens, such as Te or Se, and bulk carrier densities are reduced by orders of magnitude to achieve access to robust topological surface states without altering carrier mobility or the bandstructure.

"The main advance of this work is that the new hydrogenation process is fully reversible, as hydrogen-chalcogen moiety can be disassociated by a low-temperature annealing protocol under which hydrogen is easily removed," said Krusin-Elbaum, professor in CCNY's Division of Science. "It is also multiply-cyclable and reproducible, thereby resolving one of the key limitations of magnetic and nonmagnetic [topological](#)

[insulators](#) and can be applied not only post-growth to materials but also to fully fabricated nanodevices."

The research in the Krusin Lab centers on exploring novel quantum phenomena such as Quantum Anomalous Hall (QAH) effect, which describes an insulator that conducts dissipationless current in discrete channels on its surfaces, 2D superconductivity, and axion state phenomena featuring a quantized thermal transport, all with the potential if industrialized to advance energy-efficient technologies.

Krusin-Elbaum and her team said that the technique they have demonstrated is very general and ultimately may advance the potential of intrinsic topological magnets to transform future quantum electronics.

More information: Haiming Deng et al, Topological surface currents accessed through reversible hydrogenation of the three-dimensional bulk, *Nature Communications* (2022). [DOI: 10.1038/s41467-022-29957-3](#)

Provided by The City University of New York

Citation: Hydrogen-tuned topological insulators may lead to new platforms in sustainable quantum electronics (2022, May 4) retrieved 2 May 2024 from <https://phys.org/news/2022-05-hydrogen-tuned-topological-insulators-platforms-sustainable.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--