

Nanoscale currents improve understanding of quantum phenomena

May 5 2022



Credit: Unsplash/CC0 Public Domain

Besides charge, subatomic particles like electrons also carry a property called spin, which is responsible for magnetism. Novel proposals to use spin to store information have emerged in recent years with the promise

to be more energy efficient and to bring new functionalities to devices of communication and sensing. For his Ph.D. research, Adonai Rodrigues Da Cruz studied the spin dynamics in more detail using theory and numerical simulations. He defended his thesis at the department of Applied Physics on May 3rd.

Spin defects

The ability to control single quantum states is crucial for the development of new quantum technologies for future communication, sensing, and information processing devices. In recent years, a number of novel ways of using electron spin to store [digital information](#) in such quantum technologies have been proposed.

The use of defects in [semiconductor materials](#) carrying spin (so called spin defects) have been promoted as materials to create quantum bits—the key component of any quantum technology.

Sensors based on spin defects, such as the nitrogen-vacancy (NV) center in diamond, are already commercially available and have emerged as the most exciting new technology for nanoscale magnetic measurements. By controlling and interrogating the spin state of this crystal defect, researchers have been able to measure extremely small magnetic fields, and thus study the properties of new materials in more detail.

Understanding orbital effects

Up to now, both experimental and [theoretical studies](#) on single defects in semiconductors have focused more on the spin part and largely neglected the orbital contribution to the local properties around the defects.

For his Ph.D. research, Adonai Rodrigues Da Cruz sought a better

understand of the orbital effects. This insight was provided through the development of theoretical formalisms to describe circulating currents in different materials. Using [numerical simulations](#) and analytic expressions for the propagation of electrons in two-dimensional semiconductors, he accurately predicted how nano-currents distribute according to the surrounding environment.

Fringe magnetic fields can be generated by the circulating currents around single spin defects located inside the semiconductors. One of the main findings of Da Cruz's research is that both the magnitude and spatial dimensions of the fringe [magnetic field](#) are within the desired range of sensitivity of current NV-based probes. Therefore, a scanning NV sensor could be used as a direct probe of the internal orbital magnetism associated with single defects.

Da Cruz's work also suggests that the spatial features of the currents can be strongly tuned by external gating, opening up the possibility of electrically control short-distance [spin](#) couplings, which is essential for quantum entangling gate, an important operation in quantum computing.

More information: [Theory of nanoscale currents and effects from spin-orbit coupling in semiconductors](#)

Provided by Eindhoven University of Technology

Citation: Nanoscale currents improve understanding of quantum phenomena (2022, May 5) retrieved 24 April 2024 from <https://phys.org/news/2022-05-nanoscale-currents-quantum-phenomena.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.