

## Thin quantum wires work better with less insulating coatings

May 4 2022



The form factor for a quantum wire with finite or infinite confining potential as a function of wire radius for the ratio of dielectric constants a  $\varepsilon 2/\varepsilon 1 = 0.5$ , b  $\varepsilon 2/\varepsilon 1 = 3.0$  (solid line). Asterisks show the radius dependence of the relative difference of form factors. The linear electron density is n =  $10^6$  cm<sup>-1</sup>, the wave vector q =  $10^6$  cm<sup>-1</sup>, the temperature of the system T = 300 K. Credit: *The European Physical Journal B* (2022). DOI: 10.1140/epjb/s10051-022-00295-z



New theoretical analysis considers cases where the electrons are allowed to exist beyond the boundaries of semiconducting quantum wires—with important implications for their performance

Thin, semiconducting wires have attracted much recent attention in physics—both in experiments and <u>theoretical analysis</u>. Called quantum wires, these structures are often coated in insulating materials, and several previous studies have now explored how the mismatch between the insulating properties of both materials can influence their performance. Through new analysis published in *The European Physical Journal B*, Nguyen Nhu Dat and Nguyen Thi Thuc Hien at Duy Tan University, Vietnam, show that thinner wires with less insulating coatings can improve the mobility of the electrons they carry.

Quantum wires have a diverse array of potential applications, and their use in devices including lasers, LEDs, transistors, and sensors is now being widely explored. Results from previous studies have shown that the insulating properties of their coating materials can vary the interactions which take place between electrical charges within the semiconducting wire. In turn, this influences the quantum states of the wire's electrons. However, these models have presented conflicting conclusions about the ability of electrons to move through the wire, depending on whether coatings are more or less insulating than the semiconductor.

Through their detailed analysis, Nguyen Nhu Dat and Nguyen Thi Thuc Hien have now improved on these previous approaches. While previous studies assumed that these electrons remain completely confined to the wire, the duo considered the case where <u>electrons</u> are allowed to cross the outer boundary of the semiconductor. Their resulting calculations showed that previous models have likely underestimated electron



mobility—which becomes roughly 10 times greater in thin wires, when they are coated in less insulating materials than the semiconductor. All the same, previous models are still useful for describing thicker quantum wires. The result provides important insights into the conducting properties of quantum wires, and could enable researchers in future studies to better understand their potential applications.

**More information:** Nguyen Nhu Dat et al, Dielectric function and impurity-limited mobility of semiconductor quantum wires: effects of dielectric mismatch and finite confining potential, *The European Physical Journal B* (2022). DOI: 10.1140/epjb/s10051-022-00295-z

Provided by Springer

Citation: Thin quantum wires work better with less insulating coatings (2022, May 4) retrieved 5 May 2024 from <u>https://phys.org/news/2022-05-thin-quantum-wires-insulating-coatings.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.