


Ultrasound-guided microbubbles boost immunotherapy efficacy

May 30 2022



Credit: Unsplash/CC0 Public Domain

Researchers at The University of Texas MD Anderson Cancer Center have developed an ultrasound-guided cancer immunotherapy platform that generates systemic antitumor immunity and improves the



therapeutic efficacy of immune checkpoint blockade. The findings from the preclinical study were published today in *Nature Nanotechnology*.

As the first-of-its-kind platform, the Microbubble-assisted UltraSoundguided Immunotherapy of Cancer (MUSIC) approach employs nanocomplexes combined with microbubbles to effectively deliver cyclic guanosine monophosphate-adenosine monophosphate (cGAMP), an immunotransmitter involved in anticancer immunity, into antigenpresenting cells (APCs). Inside the APCs, the microbubbles release cGAMP to activate the GMP-AMP synthase (cGAS)-stimulator of interferon genes (STING) pathway, which stimulates type I interferon responses that are essential for priming tumor-specific T cells.

In the preclinical study, the MUSIC strategy demonstrated a complete tumor eradication rate of 60% when administered as monotherapy in breast <u>cancer</u> models. When combined with an anti-PD-1 antibody, MUSIC significantly improved antitumor responses with minimal toxicity effects, including enhanced primary tumor control and decreased systemic disease progression. In addition, the combination therapy demonstrated superior survival benefit, with a 76% increase in median survival compared to either therapy alone.

"By investigating the mechanisms of action in producing a robust STING activation, we identified a new strategy to activate both the innate and adaptive antitumor immune responses," said Wen Jiang, M.D., Ph.D., assistant professor of Radiation Oncology and the study's co-senior author. "Our findings show that the MUSIC strategy is capable of paving the way toward novel image-guided strategies for targeted cancer immunotherapy."

Immunotherapy has transformed <u>cancer treatment</u>, offering clinical benefits for patients with treatment-refractory metastatic cancers such as melanoma, <u>non-small cell lung cancer</u> and renal cell cancer. However,



not all patients respond to immune checkpoint blockade. Therefore, developing a more effective immunotherapy strategy to benefit larger numbers of cancer patients with localized and metastatic disease remains an unmet clinical need.

"Although the majority of cancer immunotherapies have focused on boosting the adaptive branch of the body's immune system, there has been a growing realization that both the innate and adaptive branches of the body's <u>immune system</u> need to be engaged to generate optimal antitumoral immunity," Jiang said. "This understanding has led to the development of new immunotherapies that target the regulators of innate immune systems, including the cGAS-STING pathway."

Natural agonists, such as cyclic dinucleotides, activate the cGAS-STING pathway, but concerns over poor cytosolic entry, serum stability and systemic toxicity have been major limitations for clinical translation. To overcome these challenges, Jiang and his collaborators developed MUSIC as the first-ever image-guided cancer immunotherapy strategy that uses antibody targeting to activate STING in APCs through delivery of molecular drugs.

Because the microbubbles also serve as <u>contrast agents</u> for ultrasound, the researchers use ultrasound scanners to image the tumor and to precisely identify the location where the microbubbles have accumulated. After confirming the microbubbles are bound to the tumors, the researchers activate ultrasonic frequencies, which cause the microbubbles to oscillate and burst, creating transient pores in the cell membrane that allow nucleic acids to be transferred directly into the cell cytosol. This technique, called sonoporation, previously has been used on tumor cells, but the MUSIC platform is the first to bind nanocomplexes to microbubbles to deliver cGAMP immunotransmitters directly into APCs.



"The beauty of our platform is that ultrasound machines are already clinically available in many outpatient settings and microbubbles are FDA-approved contrast agents for ultrasound imaging," Jiang said. "Therefore, we expect there is a very real opportunity to translate MUSIC into a clinic application to benefit cancer patients."

The same concept and design principle behind the MUSIC platform's microbubble technology could be readily translated to nanoscale systems for targeted systemic delivery and activation of innate immune sensors under image guidance for <u>cancer immunotherapy</u> applications, Jiang added.

"Our MUSIC platform is exciting because it provides a new framework for developing image-guided immunotherapy by using acoustically responsive biomaterials to enable efficient, targeted and robust immune activation to produce potent antitumor effects while minimizing systemic toxicity," Jiang said. "The versatility of the MUSIC platform could potentially be applied to targeted delivery of other immunestimulating agents, such as nucleotide-based vaccines, mRNAs and other gene therapies for multiple human diseases."

More information: Wen Jiang, Cancer immunotherapy based on image-guided STING activation by nucleotide nanocomplex-decorated ultrasound microbubbles, *Nature Nanotechnology* (2022). DOI: 10.1038/s41565-022-01134-z. www.nature.com/articles/s41565-022-01134-z

Provided by University of Texas M. D. Anderson Cancer Center

Citation: Ultrasound-guided microbubbles boost immunotherapy efficacy (2022, May 30) retrieved 26 June 2024 from <u>https://phys.org/news/2022-05-ultrasound-guided-microbubbles-</u>



boost-immunotherapy-efficacy.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.