

Validation brings new predictive capability to global megafire smoke impacts

May 17 2022



Recent megafires in Australia and British Columbia have injected unprecedented amounts of smoke into the stratosphere. Modeling led by Los Alamos National Laboratory will help predict the effects of similar future events. Credit: David Peterson, FIREX-AQ

New research modeling smoke from two recent megafires sets the stage



for better forecasting of how emissions from these global-scale events will behave and impact temperatures. As huge wildfires become more common under climate change, increased attention has focused on the intensity and duration of their emissions, which rival those of some volcano eruptions.

Megafires in British Columbia in 2017 and Australia in 2019-2020 injected massive amounts of smoke into the stratosphere, allowing firstever detailed satellite- and ground-based measurements of such cataclysms. Using that data for validation, a Los Alamos National Laboratory–led team modeled the behavior and impacts of the smoke as it rose from the lower atmosphere into the high-riding stratosphere, then circulated the globe. The research appeared in the *Journal of Geophysical Research—Atmospheres*.

"This is the only time we've tracked the global scale smoke phenomenon with satellite- and ground-based observations, which allows us to improve the <u>model</u> and understand the impact," said Manvendra Dubey, project lead and co-author of the paper published this week in the *Journal of Geophysical Research: Atmospheres.* "The models and measurements are coming together to enhance predictability."

"As fire regimes change and enter new paradigms of behavior under future <u>climate change</u>, data from past fires can't be used for prediction and assessment," said Gennaro D'Angelo, a co-author on the paper and research scientist at Los Alamos.

"The models are the only way you can forecast their smoke effects," Dubey said. "For instance, observations of the Australian fire showed that <u>black carbon</u> got a boost from <u>solar heating</u> and rose to 30 kilometers in the stratosphere, which made the plume last longer, about 16 months. Our model accounts for this self-heating phenomenon predicted by the late Robert C. Malone at Los Alamos in the 1980s—and



our new study unequivocally validates it."

Plumes have cooling effect

The 2019-2020 Australian megafire injected huge amounts of smoke and soot into the atmosphere that were observed, with global impacts on temperature as this study shows. Shading from the Australian plume lasted a few months. That effect lowered temperatures in the <u>southern</u> <u>hemisphere</u> about 0.2 degrees Celsius, information that has implications for global climate change models.

The smaller plume from the 2017 British Columbia fires did not trigger similar cooling. The study highlights when and how megafire smoke impacts <u>global climate</u>, much like volcanic sulfate and ash injections do.

More information: Gennaro D'Angelo et al, Contrasting Stratospheric Smoke Mass and Lifetime From 2017 Canadian and 2019/2020 Australian Megafires: Global Simulations and Satellite Observations, *Journal of Geophysical Research: Atmospheres* (2022). DOI: 10.1029/2021JD036249

Provided by Los Alamos National Laboratory

Citation: Validation brings new predictive capability to global megafire smoke impacts (2022, May 17) retrieved 17 May 2024 from <u>https://phys.org/news/2022-05-validation-capability-global-megafire-impacts.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.