


'Protective cloak' prevents plants from self-harming in very bright conditions

June 14 2022



Integration of environmental signals on the expression of the photoprotection-related genes. (A) Schematic summary of changing light quality and quantity throughout the day. Blue light reaches deeper levels of the water column, while red light is absorbed near the water surface. In addition, PAR can be strongly reduced by cloud cover, while UV-B radiation might even increase on partly cloudy days. While direct sunlight is shielded/reduced by canopy shading, blue light (and UV light in the case of a plant canopy) reach shaded areas more effectively than other wavelength of PAR through Rayleigh scattering, which

increases as the wavelength of light decreases. (B) Signals that regulate energy dissipation in *Chlamydomonas*. Transcription of LHCSR1, LHCSR3, and PSBS is strongly initiated with exposure to a very low amount of white light ($5 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$; Fig. 1). This activation is not only strongest for LHCSR3 but also apparent for LHCSR1 and PSBS and is dependent on the *Chlamydomonas* blue light-dependent photoreceptor PHOT1. All three transcripts are also partially regulated by PET downstream of PSII and the generation of retrograde signals by HL (red). UV-B radiation directly facilitates monomerization of the UVR8 homodimer, which then binds to COP1 and allows the participation of other factors (not included in the figure) in the transcriptional regulation of the photoprotective genes (purple). UV-B exposure may also lead to the generation of RS in the chloroplast that further triggers signaling events (red). In addition, LHCSR3 is strongly controlled by CO_2 levels and CIA5, while PSBS may be affected by CO_2 to a minor extent (orange). Additional discussion of the role of CO_2 in regulating LHCSR3 is presented in Ruiz-Sola et al. (48). The heatmap table summarizes the transcript fold change for each gene in the transition from dark to the indicated conditions. NADP⁺, nicotinamide adenine dinucleotide phosphate; NADPH, reduced form of NADP⁺; ADP, adenosine 5'-diphosphate ; ATP, adenosine 5'-triphosphate. Credit: *Science Advances* (2022). DOI: 10.1126/sciadv.abn1832

New work led by Carnegie's Petra Redekop, Emanuel Sanz-Luque, and Arthur Grossman probes the molecular and cellular mechanisms by which plants protect themselves from self-harm. Their findings, published by *Science Advances*, improve our understanding of one of the most-important biochemical processes on Earth.

Plants, [algae](#), and certain bacteria are capable of converting the sun's energy into [chemical energy](#) through a process called photosynthesis. It underpins our entire food chain and is responsible for the oxygen-rich nature of our atmosphere.

"In other words, life as we know it couldn't exist without

photosynthesis," Grossman said.

The process takes place in two stages. In the first, light is absorbed and used to synthesize energy molecules, with water a substrate and oxygen as a byproduct. These energy molecules are then used to power the second stage, in which carbon dioxide from the air is fixed into carbon-based sugars.

Photosynthesis occurs when [plants](#)—and other organisms capable of it—are exposed to the sun. But in many environments, access to sunlight can shift throughout the day due to wind, weather, clouds, and canopy cover, and other surrounding conditions.

How do [photosynthetic organisms](#) deal with this kind of dramatic variability?

"It's necessary for plants and algae to be able to harvest sufficient light when they're in the shade and to dissipate the excess of absorbed energy when light conditions are intense, such as at high noon," explained Redekop.

In extremely bright sunlight, highly reactive oxygen molecules can build up in plants and algae and cause cellular damage, or even death, if they aren't neutralized. To protect themselves from this situation, plants and algae have evolved a set of proteins that can rapidly quench the excess light energy before it can cause too much harm.

Redkop, Sanz-Luque, and Grossman, along with colleagues from CNRS and the University of Grenoble Alpes, characterized the genes that encode these protective proteins in the photosynthetic algae *Chlamydomonas* and probed the organism's ability to regulate their expression.

"We found that activation of these genes is controlled by an integrated series of signals with built-in redundancies that help plants and algae to optimize their environmental responses depending on the time of day and the surrounding environment," Grossman said.

The team's work focused on the genes that encode three *Chlamydomonas* proteins known to play a major role in this so-called photoprotection process, revealing a suite of conditions that triggered the genes' activation.

The researchers found that expression of genes encoding these proteins accumulated even in very low levels of light. This is evidence that algae and plants marshal their resources at dawn in preparation for the first few hours of intense morning light.

What's more, the activation of the genes encoding these proteins was induced by the detection of blue light, which increases from dawn to midday and decreases from noon to sunset, especially in aquatic environments. This demonstrates that the system regulating photoprotection is effective in a variety of ecosystems.

The [genes](#) are also activated by the presence of UV-B radiation, which is not blocked by [cloud cover](#), allowing algae and plants to track the time of day and prepare for the accompanying changes in light availability, even in low-[light conditions](#).

Last, one of the photoprotection proteins was regulated by the availability of carbon dioxide. The researchers say further analysis is needed to understand this integrated regulatory network.

"Taken together, this set of regulatory features form a protective cloak that dampens the risk posed by excess light in a rapidly changing environmental landscape," concluded Redekop. "This work reveals the

stunning array of mechanisms plants and algae have evolved to maintain productivity and minimize harm."

More information: Petra Redekop et al, Transcriptional regulation of photoprotection in dark-to-light transition—More than just a matter of excess light energy, *Science Advances* (2022). [DOI: 10.1126/sciadv.abn1832](https://doi.org/10.1126/sciadv.abn1832)

Provided by Carnegie Institution for Science

Citation: 'Protective cloak' prevents plants from self-harming in very bright conditions (2022, June 14) retrieved 26 March 2023 from <https://phys.org/news/2022-06-cloak-self-harming-bright-conditions.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.