


# With cryo-EM, scientists determine structure of key factor in RNA quality control

June 9 2022



Graphical abstract. Credit: *Cell* (2022). DOI: 10.1016/j.cell.2022.04.016

In biology, getting rid of stuff can be just as important as making it. A buildup of cells, proteins, or other molecules that are no longer needed can cause problems, so living things have evolved several ways to clean house.

A prime example is the RNA exosome. RNA molecules perform many roles in cells. Some of them are translated into proteins; others form a cell's protein-building machinery. The RNA exosome is a cellular machine that degrades RNA molecules that are faulty, harmful, or no longer needed. Without this microscopic Marie Kondo to prune what doesn't spark joy, our cells would become dysfunctional hoarders, unable to function.

"RNA surveillance and degradation pathways exist in all forms of life," explains Christopher Lima, Chair of the Structural Biology Program in the Sloan Kettering Institute. "From bacteria to humans, all living things have mechanisms to monitor the quality of RNA and to purposely degrade it."

For a long time, Dr. Lima says, these pathways were considered, like housework, kind of boring. But it turns out that these degradation pathways are highly regulated and control everything from embryonic development to the progression of the cell cycle.

What's more, errors in these pathways can lead to many types of disease, from cancer to neurodegeneration.

In a new paper published June 9, 2022 in *Cell*, Dr. Lima and M. Rhyan Puno, a postdoctoral fellow in the Lima lab, present findings that help

explain how the RNA exosome locates the RNA that needs to be degraded. With the help of cryogenic-electron microscopy (cryo-EM), an advanced type of imaging technology, the scientists were able to decipher the structure of a protein assembly called Nuclear Exosome Targeting (NEXT) Complex, which is a key part of the degradation machinery.

"We knew that NEXT targets and delivers RNA to the exosome, but biochemically and structurally, we didn't have a clue what it looks like or how it works," Dr. Puno says.

Now, with cryo-EM, the scientists have obtained the first clear pictures of NEXT bound to RNA. These pictures, in conjunction with accompanying biochemical and biological experiments, offer hints about how RNA molecules are handed over to the exosome for destruction.

## **Inching closer to a structure**

Several years ago, Dr. Puno began studying NEXT's structure using the then gold standard approach of X-ray crystallography. In this method, proteins are first made into crystals, with the proteins all aligned in the same way. Then, X-rays are passed through the crystals and the pattern of X-rays that hit a detector can be interpreted to determine the protein's structure.

While Dr. Puno was able to crystallize the NEXT protein, the resulting X-ray diffraction pictures weren't good enough to see details of the structure.

"But then came the cryo-EM revolution," he says. "Cryo-EM helped us visualize what this protein looks like and how it binds its RNA substrates."

## Visualizing proteins in motion

Cryo-EM works by capturing many different images of a frozen but non-crystallized sample of a protein, and then using computational methods to align them into a finalized sharp image.

"It's almost like capturing a bunch of pictures of a bird in flight," Dr. Lima says. "There's all sorts of confusing movements and the bird's wings can look blurry. But if we can find parts of the wing in all those different pictures, then we can align the pictures to reconstruct what the wings of the bird look like and determine how they work."

From the cryo-EM pictures, the scientists were able to see that the NEXT proteins form a very flexible dimer—meaning that two copies of NEXT proteins join together as a functional unit.

"That was really, really puzzling," Dr. Puno says, noting that dimer formation hasn't been visualized before for these types of proteins.

"From biochemical experiments we performed, we know that dimerization is somehow important for degradation," he continues. "But it's still a mystery to us what role the dimer plays in guiding RNA to the exosome."

To help solve the mystery, they hope to capture the NEXT complex interacting at different steps in the degradation process and then visualize these conformations with cryo-EM.

## RNA degradation and disease

There are big stakes involved. An indication of just how important RNA degradation is comes from the long list of diseases that result from

defective or poorly controlled degradation. Perhaps the most famous example is cystic fibrosis. In this case, the messenger RNA encoding a protein that shuttles ions across cell membranes is degraded by RNA decay pathways. As a result, the protein is not present in mucous membranes of the lungs, which leads to a buildup of mucus there and results in severely impaired breathing.

"It's a famous example of RNA quality control with bad results," Dr. Lima says.

But defects in RNA degradation pathways also play a role in several types of cancer. In fact, two of the [genetic mutations](#) that MSK's genetic testing platform, MSK-IMPACT, tests for are found in genes related to the RNA exosome pathway, including a [protein](#) in NEXT.

And it's not only messenger RNA that needs proper quality control, Dr. Lima explains.

"The reality is if you have defective RNA quality-control pathways, your ribosomes don't work, your transfer RNAs don't work, your spliceosomes don't work." The list goes on and on.

The breadth of functions that RNA performs explains why defective RNA degradation pathways can have such cascading disease-causing effects.

Making sense of these effects will require a deeper and more extensive understanding of not just the RNA exosome itself, but also the "upstream" proteins, like NEXT, that help surveil RNA and decide when an RNA is defective or no longer needed.

"The dream is to initiate the RNA degradation reaction, put the sample in the cryo-EM, and actually see all the possible confirmations as it's

doing its job," Dr. Lima says. "As structural biologists, we want to be able to see processes in action and then be able to reassemble them."

**More information:** M. Rhyan Puno et al, Structural basis for RNA surveillance by the human nuclear exosome targeting (NEXT) complex, *Cell* (2022). [DOI: 10.1016/j.cell.2022.04.016](https://doi.org/10.1016/j.cell.2022.04.016)

Provided by Memorial Sloan Kettering Cancer Center

Citation: With cryo-EM, scientists determine structure of key factor in RNA quality control (2022, June 9) retrieved 24 May 2024 from <https://phys.org/news/2022-06-cryo-em-scientists-key-factor-rna.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--