

Dynamic rearrangement and autophagic degradation of the mitochondria during plant spermiogenesis

June 16 2022



During spermiogenesis, the mitochondrial number is reduced to one through fission and autophagic degradation, and subsequently increased to two through fission of the anterior mitochondrion. Other cytoplasmic components are later degraded through autophagic (and non-autophagic) degradation. Credit: NIBB

A collaborative team of researchers from the National Institute for Basic Biology, the University of Tokyo, and Gunma University in Japan reported that in the liverwort Marchantia polymorpha, the number of



mitochondria in the spermatozoid (sperm) is controlled by autophagy during spermiogenesis.

Most <u>eukaryotic cells</u> possess <u>mitochondria</u> in greatly varying numbers and shapes depending on the cell type and cellular conditions. Compared with other <u>cell types</u>, eukaryotic sperm cells generally exhibit a characteristic mitochondrial morphology and distribution, while also exhibiting a remarkable diversity among species.

Though the word "sperm" brings to the mind images of the mammalian sperms that "swim," certain groups of plants also use sperms for sexual reproduction. One such group is the Bryophytes that are commonly found in city parks and backyards of homes.

A single bryophyte <u>somatic cell</u> possesses anywhere from tens to hundreds of mitochondria. By contrast, the spermatozoid cell possesses a fixed number of two mitochondria: one in the head and the other in the tail of the cell body. However, the mechanism by which these two mitochondria are formed during spermiogenesis remains unclear.

In an article published in *Cell Reports*, this research team examined in detail how the number and shape of mitochondria change during the transformation of spermatids into spermatozoids (spermiogenesis) in liverworts.





The wild-type spermatozoids possess two mitochondria, whereas, most defective spermatozoids in mitochondrial fission possess only one anterior mitochondrion. The mitochondria are shown in green. Credit: NIBB

Dr. Takuya Norizuki, the first author of the article, spoke of the team's findings that in liverwort, the number of mitochondria decreases as spermiogenesis progresses until only one remains. The remaining mitochondrion then divides asymmetrically into two mitochondria; the larger mitochondrion becomes the anterior mitochondrion and the smaller one becomes the posterior mitochondrion. When this mitochondrial division was inhibited, sperms with only an anterior mitochondrion were formed.

The research team investigated the mechanism underlying the mitochondrial number reduction, with a special focus on autophagy. Autophagy degrades cellular components, including the organelles, by engulfing them into double-membraned sacs (autophagosomes) that fuse with the vacuoles/lysosomes.



When the functions of the genes required for autophagy were lost, the mitochondria were not degraded during spermiogenesis, and a large number of mitochondria remained in the mutant spermatozoid. The research team also found through <u>electron microscopy</u> that mitochondria were selectively engulfed by autophagosomes during spermiogenesis in the wild-type liverworts. These results indicated that autophagy reduces the number of mitochondria during spermiogenesis.



Autophagy-defective spermatozoids possess a larger number of mitochondria, as opposed to two mitochondria in wild-type spermatozoids. Mitochondria and nuclei are colored green and blue, respectively. Credit: NIBB

Professor Takashi Ueda, the leader of the research team, commented that "this study reveals how the characteristic mitochondrial pattern in bryophyte spermatozoids arises and that autophagy is deeply involved in this process. During the process of spermiogenesis in animals, the



removal of excessive cytoplasm occurs, but <u>autophagy</u> does not seem to play a substantial role in this process. Instead, a neighboring cell 'takesup' the excessive cytoplasm and organelles by phagocytosis to remove it from the developing <u>sperm</u>. Thus, plants and animals have employed completely different mechanisms to remove organelles during spermiogenesis."

More information: Takuya Norizuki et al, Dynamic rearrangement and autophagic degradation of mitochondria during spermiogenesis in the liverwort Marchantia polymorpha, *Cell Reports* (2022). <u>DOI:</u> <u>10.1016/j.celrep.2022.110975</u>

Provided by National Institutes of Natural Sciences

Citation: Dynamic rearrangement and autophagic degradation of the mitochondria during plant spermiogenesis (2022, June 16) retrieved 6 May 2024 from https://phys.org/news/2022-06-dynamic-rearrangement-autophagic-degradationmitochondria.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.