

## **Global water basins hotspots prioritize areas under threat**

June 29 2022, by Mark Ferguson



Global co-occurrence of freshwater stress and storage trends. **a** Freshwater stress, derived from freshwater withdrawal and streamflow datasets. **b** Freshwater storage trend per basin. **c** Combinations of freshwater stress and storage trend per basin, which together derive basin freshwater status. Values overlaying the legend indicate the number of basins satisfying each set of conditions. For categorical plotting purposes only,  $\pm 3$  mm year<sup>-1</sup> is used as the threshold denoting a clear directional storage trend, based on the error level of the underlying observations. **d**-**g** The exposure of social-ecological activity to



freshwater stress and storage trends. Each plot represents storage trends as the *x* -axis coordinate, and log-transformed freshwater stress as the y-axis coordinate with the size of each circle based on the basin's value respective to each plotting dimension. Credit: *Nature Communications* (2022). DOI: 10.1038/s41467-022-28029-w

New research at the intersection of how humans and ecosystems interact with water shows that the most-stressed regions in the world are becoming drier, leading to water governance, economic and social challenges.

Recent studies involving a University of Saskatchewan (USask) researcher show where groundwater is becoming less available globally, but less work has been done to understand the impact changes to these <u>water systems</u> have on people. That is why a new study, which was recently published in *Nature Communications*, takes into account both social and ecological impacts, allowing researchers to identify global "hotspots" where the threat to the water <u>basin</u> is most concerning.

"There's a consensus in <u>sustainability science</u> and sustainability literature around being wary of broad-brush solutions," said Xander Huggins, a Ph.D. candidate with USask's Global Institute for Water Security (GIWS), and the University of Victoria Department of Civil Engineering.

"These [solutions] really need to be generated locally, and they will be different from each other from place to place... This gives us a template for where to focus globally with greater specificity."

Huggins, the lead author of the paper, says this is "a call to action," and there's already been interest in the research from non-governmental



organizations such as the World Wildlife Fund.

Once the vulnerable water basins are clearly identified, place-based solutions can be implemented by <u>local communities</u>, organizations and governments.

"There is an urgent need to address these drying water basins. The most vulnerable basins impact more than 1.5 billion people, 17% of global food crop production, 13% of global gross domestic product, and hundreds of significant wetlands," said Huggins.

He said one of the ways governments and communities can address this crisis is to apply integrated water resources management (IWRM), a process that promotes co-ordinated water management and is tied into the United Nations' Sustainable Development Goals.

The study suggests that basins with low levels of IWRM where vulnerability is high should be the priority basins. Yet comparing the vulnerability mapping they produced to the current level of IWRM implementation, they found that transboundary basins—shared by multiple countries—are the least likely to implement such measures and are also the most at risk. Nations with low levels of IWRM implementation and very high vulnerability include Afghanistan, Algeria, Argentina, Egypt, India, Iraq, Kazakhstan, Mexico, Somalia, Ukraine, Uzbekistan, and Yemen.

"Part of the reason why they are so vulnerable is that there's less diplomacy... between these multi-jurisdictional water systems that are affecting humans and ecosystems at once," Huggins said.

The paper calls for greater policy integration and hydro-diplomacy, noting that of the nearly 700 water conflicts documented since 2000 by The Water Conflict Chronology, two-thirds (68%) are found within



either transitional or hotspot basins.

Jay Famiglietti, executive director of GIWS and co-supervisor for Huggins's Ph.D. work, says the research helps prioritize the attention that these different basins need.

"The work we've done until this point has been strictly looking at water availability and changes in water availability, and we've only talked in broad strokes about how it might impact food production, how it might impact biodiversity, what it means for political stability or conflict," Famiglietti said.

"This is really the first step at putting a few of those other stressors together, along with changing <u>water availability</u>, to get a more robust picture of just how vulnerable these basins are."

Looking to the future, Famiglietti, Huggins and others are putting together an international, transdisciplinary group to address the problem of global groundwater sustainability, and Famiglietti said research like this is critical to moving forward.

"You really have to understand the <u>social structure</u>, the governance structure, the biodiversity that's at risk, the wealth of the nation, its capacity," he said. "This is really the first step towards that diagnosis."

**More information:** Xander Huggins et al, Hotspots for social and ecological impacts from freshwater stress and storage loss, *Nature Communications* (2022). DOI: 10.1038/s41467-022-28029-w

Provided by University of Saskatchewan



Citation: Global water basins hotspots prioritize areas under threat (2022, June 29) retrieved 27 June 2024 from <u>https://phys.org/news/2022-06-global-basins-hotspots-prioritize-areas.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.