

## Found: The 'holy grail of catalysis'—turning methane into methanol under ambient conditions using light

June 30 2022



Credit: ORNL/Jill Hemman

An international team of researchers, led by scientists at the University



of Manchester, has developed a fast and economical method of converting methane, or natural gas, into liquid methanol at ambient temperature and pressure. The method takes place under continuous flow over a photo-catalytic material using visible light to drive the conversion.

To help observe how the process works and how selective it is, the researchers used neutron scattering at the VISION instrument at Oak Ridge National Laboratory's Spallation Neutron Source.

The method involves a continuous flow of methane/oxygen-saturated water over a novel metal-organic framework (MOF) catalyst. The MOF is porous and contains different components that each have a role in absorbing light, transferring electrons and activating and bringing together methane and oxygen. The liquid methanol is easily extracted from the water. Such a process has commonly been considered "a holy grail of catalysis" and is an area of focus for research supported by the U.S. Department of Energy. Details of the team's findings, titled "Direct photo-oxidation of methane to methanol over a mono-iron hydroxyl site," are published in *Nature Materials*.

Naturally occurring methane is an abundant and valuable fuel, used for ovens, furnaces, water heaters, kilns, automobiles and turbines. However, methane can also be dangerous due to the difficulty of extracting, transporting and storing it.

Methane gas is also harmful to the environment when it is released or leaks into the atmosphere, where it is a potent greenhouse gas. Leading sources of atmospheric methane include fossil fuel production and use, rotting or burning biomass such as forest fires, agricultural waste products, landfills and melting permafrost.

Excess methane is commonly burned off, or flared, to reduce its



environmental impact. However, this combustion process produces carbon dioxide, which itself is a greenhouse gas.

Industry has long sought an economical and efficient way to convert methane into methanol, a highly marketable and versatile feedstock used to make a variety of consumer and industrial products. This would not only help reduce methane emissions, but it would also provide an economic incentive to do so.

Methanol is a more versatile carbon source than methane and is a readily transportable liquid. It can be used to make thousands of products such as solvents, antifreeze and acrylic plastics; synthetic fabrics and fibers; adhesives, paint and plywood; and chemical agents used in pharmaceuticals and agrichemicals. The conversion of methane into a high-value fuel such as methanol is also becoming more attractive as petroleum reserves dwindle.

## Breaking the bond

A primary challenge of converting methane  $(CH_4)$  to methanol  $(CH_3OH)$  has been the difficulty of weakening or breaking the carbon-hydrogen (C-H) <u>chemical bond</u> in order to insert an oxygen (O) atom to form a C-OH bond. Conventional methane conversion methods typically involve two stages, steam reforming followed by syngas oxidation, which are energy intensive, costly and inefficient as they require high temperatures and pressures.

The fast and economical methane-to-methanol process developed by the research team uses a multicomponent MOF material and <u>visible light</u> to drive the conversion. A flow of  $CH_4$  and  $O_2$  saturated water is passed through a layer of the MOF granules while exposed to the light. The MOF contains different designed components that are located and held in fixed positions within the porous superstructure. They work together



to absorb light to generate electrons which are passed to oxygen and methane within the pores to form methanol.

"To greatly simplify the process, when methane gas is exposed to the functional MOF material containing mono-iron-hydroxyl sites, the activated oxygen molecules and energy from the light promote the activation of the C-H bond in methane to form methanol," said Sihai Yang, a professor of chemistry at Manchester and corresponding author. "The process is 100% selective—meaning there is no undesirable byproduct—comparable with methane monooxygenase, which is the enzyme in nature for this process."

The experiments demonstrated that the solid catalyst can be isolated, washed, dried and reused for at least 10 cycles, or approximately 200 hours of reaction time, without any loss of performance.

The new photocatalytic process is analogous to how plants convert light energy to chemical energy during photosynthesis. Plants absorb sunlight and carbon dioxide through their leaves. A photocatalytic process then converts these elements into sugars, oxygen and water vapor.

"This process has been termed the 'holy grail of catalysis.' Instead of burning methane, it may now be possible to convert the gas directly to methanol, a high-value chemical that can be used to produce biofuels, solvents, pesticides and fuel additives for vehicles," said Martin Schröder, vice president and dean of faculty of science and engineering at Manchester and corresponding author. "This new MOF material may also be capable of facilitating other types of chemical reactions by serving as a sort of test tube in which we can combine different substances to see how they react."

## Using neutrons to picture the process



"Using neutron scattering to take 'pictures' at the VISION instrument initially confirmed the <u>strong interactions</u> between  $CH_4$  and the monoiron-hydroxyl sites in the MOF that weaken the C-H bonds," said Yongqiang Cheng, instrument scientist at the ORNL Neutron Sciences Directorate.

"VISION is a high-throughput neutron vibrational spectrometer optimized to provide information about molecular structure, chemical bonding and intermolecular interactions," said Anibal "Timmy" Ramirez Cuesta, who leads the Chemical Spectroscopy Group at SNS. "Methane molecules produce strong and characteristic neutron scattering signals from their rotation and vibration, which are also sensitive to the local environment. This enables us to reveal unambiguously the bondweakening interactions between  $CH_4$  and the MOF with advanced neutron spectroscopy techniques."

## Fast, economical and reusable

By eliminating the need for high temperatures or pressures, and using the energy from sunlight to drive the photo-oxidation process, the new conversion method could substantially lower equipment and operating costs. The higher speed of the process and its ability to convert <u>methane</u> to methanol with no undesirable byproducts will facilitate the development of in-line processing that minimizes costs.

**More information:** Sihai Yang, Direct photo-oxidation of methane to methanol over a mono-iron hydroxyl site, *Nature Materials* (2022). DOI: 10.1038/s41563-022-01279-1. www.nature.com/articles/s41563-022-01279-1

Provided by Oak Ridge National Laboratory



Citation: Found: The 'holy grail of catalysis'—turning methane into methanol under ambient conditions using light (2022, June 30) retrieved 1 May 2024 from <a href="https://phys.org/news/2022-06-holy-grail-catalysisturning-methane-methanol.html">https://phys.org/news/2022-06-holy-grail-catalysisturning-methane-methanol.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.