

# New prehistoric plant discovery highlights the unexpected diversity of non-flowering plants

June 14 2022, by Az (Ashley A.) Klymiuk

---



Holberg Inlet, along the north shore of Vancouver Island, is a rich site for Cretaceous flora. Credit: Az Klymiuk, Author provided

The renowned Apple Bay fossil locality of northern Vancouver Island is helping us reimagine seed plant diversity in the Early Cretaceous, the last of three geologic periods comprising the Mesozoic Era.

Until recently, it was assumed that most major lineages of seed plants

had been recognized in the fossil record. Their evolution and relationships to one another are still being debated, but few botanists believed that we might still be missing wide swathes of vascular plant diversity.

Using fossils from this site, my colleagues (paleobotanists Gar Rothwell and Ruth Stockey) and I have been challenging the assumption that the plant fossil record is largely complete. Our recent discovery of [a new genus of non-flowering seed plants](#) illustrates that researchers' understanding of seed plant diversity in the Cretaceous is by no means complete.

## **Exceptionally detailed fossilization**

In Apple Bay, sandstone and siltstone beds tilt gently up, variably exposed with every falling tide, and giving few clues as to their scientific significance. This site is a [konservat lagerstätte](#): a fossil locality with exceptional preservation.


During the Early Cretaceous (about 136 million years ago), seasonal storms and flooding events swept vegetation out into a marine basin. Plant debris rapidly settled and fossilized at a cellular level of detail. This type of fossilization is called [permineralization](#).

Permineralized plants offer a wealth of anatomical information, enabling us to ascertain features of plant development, organography (the arrangement of plant organs) and reproduction—data critical for understanding plant evolution. In the two decades we've been studying Apple Bay, it has become apparent that this site contains the most comprehensive record of terrestrial vegetation immediately prior to the evolution and radiation of [angiosperms \(flowering plants\)](#).

## Flower power

A mere five to 10 million years after the Apple Bay flora settled into marine sediments, [angiosperms were becoming abundant in the fossil record](#). Through the Late Cretaceous, they came to dominate terrestrial landscapes.

Prevailing narratives in [plant biology](#) suggest angiosperms out-competed other seed plants. Perhaps angiosperms were [more adaptive to changing climate](#); used nutrients [more effectively](#); or gymnosperms—non-flowering seed plants—[were already in decline by the Early Cretaceous](#), enabling angiosperms to opportunistically occupy newly available niche space.



A fossilized seed and cupule of *Xadzigacalix quatsinoensis*. Credit: Az Klymiuk, Author provided

The vast majority of seed plants that have ever lived are now extinct. The gymnosperms that are still alive are themselves relics of a much greater fossil diversity, and many major lineages of seed plants are known only from the [fossil record](#). The permineralized plants of Apple Bay demonstrate that seed plants were even more diverse throughout the Cretaceous than botanists had previously realized.

Gymnosperms are well-represented in the Apple Bay flora. They include [conifers](#) and [gnetophytes](#), which still have living exemplars like *Welwitschia*. They also include [bennettitalean](#) and [doylealean](#) plants, which represent wholly extinct orders. Indeed, the extinct Doyleales were [first described from Apple Bay in 2009](#), and have subsequently been [found in Mongolia as well](#).

## **Plant discovery**

The new plant, *Xadzigacalix quatsinoensis*, probably also represents yet another order of extinct plants. The name is a nod to Kwakwaka-speaking First Nations—the Apple Bay locality occurs within traditional and unceded territory of the Quatsino First Nation. *Xadzigacalix* derives from *xadziga*, a Kwakwaka word for plant resin, and *calix*, which is Latin for chalice. The name reflects the resinous, fleshy cupule that surrounds and protected the seeds of the *Xadzigacalix* plant.

The seeds of this plant have complex three-layered seed coats and were produced at the tip of a woody stem. A cupule encloses most of the seed,

except for a narrow tube through which the embryo would have been pollinated. This organography, along with [anatomical features](#) like vascular tissue that are observable at a cellular level, constitute a novel combination of features not found in any other lineage of plants.

It's not yet clear which group of seed plants actually gave rise to the angiosperms. Xadzigacalix likely represents a new order of seed plants, perhaps distantly related to gnetophytes or angiosperms. While research in plant molecular systematics (a discipline that uses [genetic data](#) to understand plant relationships) has resolved most [evolutionary relationships](#) within flowering plants, their origins and nearest relatives remain mysterious.

Part of this problem is that the group of seed plants most closely related to angiosperms is almost certainly extinct. Several major groups of [cupulate Mesozoic gymnosperms](#), sometimes called corystosperms or seed ferns, have been suggested as possible relatives of angiosperms. The Xadzigacalix plant is yet another possible contender to the broader "family tree" of flowering plants.

## **Perplexing origins**

Our assessment of the evolutionary context of the Xadzigacalix plant led us to re-evaluate the other groups of plants frequently [suggested as angiosperm relatives](#). Throughout the Mesozoic, many seed plants "experimented" with an array of different seed-enclosing structures.

We examined whether these cupulate structures are developmentally derived from leaf or shoot tissues. Across all Mesozoic seed plants, cupules appear to be a case of evolutionary convergence, where different [seed plants](#) independently evolved similar features. It's by no means clear if the cupules of any group are genuinely similar to the carpels—the female organs—of flowering plants.

I believe the reason [angiosperm](#) origins remain perplexing is that we still stand to discover the entirety of Mesozoic plant diversity. Our discovery of Xadzigacalix hints at how much there is left to learn.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: New prehistoric plant discovery highlights the unexpected diversity of non-flowering plants (2022, June 14) retrieved 28 June 2024 from <https://phys.org/news/2022-06-prehistoric-discovery-highlights-unexpected-diversity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.