

Real-time imaging of dynamic atom-atom interactions

June 14 2022

Imaging Dynamic Bond Formation at Atomic Resolution

Observation and However, conventional characterization of dynamic bonding methods are inadequate for predicted between capturing the atoms can open up bond formation new research frontiers dynamics Visualizing labile dimers/trimers with annular dark field scanning transmission electron microscopy (ADF-STEM) Au Time-lapsed STEM image 0.26 nm 0.25 nm Cu Ag 0.24 nm 0.244 nm 0.248 nm Ag Au Au Au Isolated metallic atoms Sequential imaging Homo/Hetero-metallic dimers and deposited on graphene via -----> and characterization with trimers (Au-Au, Au-Ag, and Au-Ag-Cu) arc plasma-deposition ADF-STEM + video tracking are observed A combination of ADF-STEM and video recording can be used

to capture bond formation dynamics at the atomic scale

Dynamic hetero-metallic bondings visualized by sequential atom imaging

Inazu et al. (2022) | Nature Communications | DOI: 10.1038/s41467-022-30533-y





Credit: Tokyo Institute of Technology

In a breakthrough Tokyo Tech researchers have managed to observe and characterize dynamic assembly of metallic atoms using an ingenious combination of scanning transmission electron microscopy and a videobased tracking. By visualizing short-lived molecules, such as metallic dimers and trimers, that cannot be observed using traditional methods, the researchers open up the possibility of observing more such dynamic structures predicted by simulations.

Chemistry is the study of bond formation (or dissociation) between atoms. The knowledge of how chemical bonds form is, in fact, fundamental to not just all of chemistry but also fields like materials science. However, traditional chemistry has been largely limited to the study of stable compounds. The study of dynamic assembly between atoms during a chemical reaction has received little attention. With recent advances in <u>computational chemistry</u>, however, dynamic, shortlived structures are gaining importance. Experimental observation and characterization of dynamic bonding predicted between atoms, such as the formation of metallic dimers, could open up new research frontiers in chemistry and materials science.

However, observing this bond dynamics also requires the development of a new methodology. This is because conventional characterization techniques only provide a time-averaged structural information and are, thus, inadequate for observing the bonds as they are formed.

Against this backdrop, researchers from Japan led by Associate Professor Takane Imaoka from Tokyo Institute of Technology (Tokyo Tech) has now provided an ingenious solution. In their study published in *Nature Communications*, the team used a combination of video



tracking and a technique called "annular dark field scanning <u>transmission</u> <u>electron microscopy</u>" (ADF-STEM) to perform sequential imaging of different metal atoms interacting with one another. This allowed them to directly observe transient structures resulting from an assembly of two similar atoms (homo-metallic dimers), two different atoms (hetero-metallic dimers), and three different atoms (hetero-metallic trimers).

The team began by depositing atoms of atoms of gold (Ag), silver (Ag), and copper (Cu) on a graphene nanoplate using a method called "arc-plasma deposition." To ensure that enough isolated single atoms were available, the deposition was limited to approximately 0.05–0.015 monolayers and high-magnification observations were performed on the flat regions of the graphene substrate.

"The elemental identification of the atoms was available with real-time tracking of the moving atoms, while ADF-STEM allowed the <u>atoms</u> to be observed under electron dose. This helped us avoid high current densities typically needed for single-atom analysis, which can cause material damage," explains Dr. Imaoka.

Additionally, ADF-STEM imaging showed an extremely high atom discrimination accuracy, ranging from 98.7% for Au–Ag to 99.9% for Au–Cu pairs. Other pairings also showed similar levels of discrimination. Moreover, the team was also able to observe Au–Ag–Cu, an extremely short-lived hetero-metallic trimer.

"Although our snapshots did not perfectly agree with the structures predicted by theoretical calculations, the average bond lengths between the elements in the observed structures are in good agreement with the calculations," says Dr. Imaoka.

The findings of this study could lead to rapid developments in nanoscience, where the characterization of metal clusters and



subnanoparticles is gaining importance, and, in the process, open doors to a completely new realm of matter.

More information: Minori Inazu et al, Dynamic hetero-metallic bondings visualized by sequential atom imaging, *Nature Communications* (2022). DOI: 10.1038/s41467-022-30533-y

Provided by Tokyo Institute of Technology

Citation: Real-time imaging of dynamic atom-atom interactions (2022, June 14) retrieved 17 July 2024 from <u>https://phys.org/news/2022-06-real-time-imaging-dynamic-atom-atom-interactions.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.