

Projected increase in space travel may damage ozone layer

June 22 2022



This graph depicts the yearly global stratospheric black carbon burden generated by 10,000 metric tons of soot exhaust per yer (black), 30,000 metric tons per year (blue), and 100,000 metric tons per year (green). The 50-year time series



average is labeled directly above each line. The red shading represents the first 10 years of the simulation that are excluded from the rest of the analysis. Current spaceflight activity contribute about 1,000 tons of black carbon to the stratosphere every year. Credit: NOAA Chemical Sciences Laboratory

Projected growth in rocket launches for space tourism, moon landings, and perhaps travel to Mars has many dreaming of a new era of space exploration. But a NOAA study suggests that a significant boost in spaceflight activity may damage the protective ozone layer on the one planet where we live.

Kerosene-burning <u>rocket engines</u> widely used by the global launch industry emit exhaust containing black carbon, or <u>soot</u>, directly into the stratosphere, where a layer of <u>ozone</u> protects all living things on the Earth from the harmful impacts of ultraviolet radiation, which include skin cancer and weakened immune systems in humans, as well as disruptions to agriculture and ecosystems.

According to new NOAA research published in the *Journal of Geophysical Research: Atmospheres*, a 10-fold increase in hydrocarbon fueled launches, which is plausible within the next two decades based on recent trends in space traffic growth, would damage the <u>ozone layer</u>, and change <u>atmospheric circulation patterns</u>.

"We need to learn more about the potential impact of hydrocarbonburning engines on the stratosphere and on the climate at the surface of the Earth," said lead author Christopher Maloney, a CIRES research scientist working in NOAA's Chemical Sciences Laboratory. "With further research, we should be able to better understand the relative impacts of different rocket types on climate and ozone."



Launch rates have tripled

Launch rates have more than tripled in recent decades, Maloney said, and accelerated growth is anticipated in the coming decades. Rockets are the only direct source of human-produced aerosol pollution above the troposphere, the lowest region of the atmosphere, which extends to a height of about four to six miles above the Earth's surface.

The research team used a climate model to simulate the impact of approximately 10,000 metric tons of soot pollution injected into the stratosphere over the northern hemisphere every year for 50 years. Currently, an estimated 1,000 tons of rocket soot exhaust are emitted annually. The researchers caution that the exact amounts of soot emitted by the different hydrocarbon fueled engines used around the globe are poorly understood.

The researchers found that this level of activity would increase annual temperatures in the stratosphere by $0.5-2^{\circ}$ Celsius (or approximately 1–4°Farenheit), which would change global circulation patterns by slowing the subtropical jet streams as much as 3.5%, and weakening the stratospheric overturning circulation.

How rocket exhaust affects the ozone layer

Stratospheric ozone is strongly influenced by temperature and atmospheric circulation, noted co-author Robert Portmann, a research physicist with the Chemical Sciences Laboratory, so it was no surprise to the research team that the model found changes in stratospheric temperatures and winds also caused changes in the abundance of ozone. The scientists found ozone reductions occurred poleward of 30 degrees North, or roughly the latitude of Houston, in nearly all months of the year. The maximum reduction of 4% occurred at the North Pole in June.



All other locations north of 30° N experienced at least some reduced ozone throughout the year. This spatial pattern of ozone loss directly coincides with the modeled distribution of black carbon and the warming associated with it, Maloney said.

"The bottom line is projected increases in rocket launches could expose people in the Northern Hemisphere to increased harmful UV radiation," Maloney said.

The research team also simulated two larger emission scenarios of 30,000 and 100,000 tons of soot pollution per year to better understand the impacts of an extremely large increase in future space travel using hydrocarbon-fueled engines, and more clearly investigate the feedbacks that determine the atmosphere's response. Results showed that the stratosphere is sensitive to relatively modest black carbon injections. The larger emission simulations showed a similar, yet more severe disruptions of atmospheric circulation and climate loss than the 10,000 metric ton case.

Building a research foundation

The study built on previous research by members of the author team. <u>A</u> 2010 study led by co-author Martin Ross, a scientist with The Aerospace Corporation, first explored the climate impact of an increase in soot-producing <u>rocket launches</u>. A second study performed at <u>NOAA in 2017</u>, on which Ross was a co-author, examined the climate response to water vapor emissions from a proposed reusable space launch system utilizing cleaner hydrogen-fueled rockets.

"Our work emphasizes the importance of ozone depletion caused by soot particles emitted by liquid-fueled rockets," Ross said. "These simulations change the long-held belief that spaceflight's only threat to the ozone layer was from solid-fueled rockets. We've shown that particles are



where the action is for spaceflight's impacts."

While the new research describes the influence that soot in rocket exhaust has on the climate and composition of the stratosphere, the scientists said it represents an initial step in understanding the spectrum of impacts on the stratosphere from increased space flight.

Combustion emissions from the different <u>rocket</u> types will need to be evaluated, they said. Soot and other particles generated by satellites burning up when they fall out of orbit is also a growing, poorly understood source of emissions in the middle-to-upper atmosphere. These and other topics will need further research to produce a complete picture of space industry emissions and their impacts on Earth's climate and ozone.

More information: Christopher M Maloney et al, The Climate and Ozone Impacts of Black Carbon Emissions From Global Rocket Launches, *Journal of Geophysical Research: Atmospheres* (2022). DOI: 10.1029/2021JD036373

Provided by NOAA Headquarters

Citation: Projected increase in space travel may damage ozone layer (2022, June 22) retrieved 23 May 2024 from <u>https://phys.org/news/2022-06-space-ozone-layer.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.