


Webb's mid-infrared spectroscopy will reveal molecules, elements

June 17 2022, by Thaddeus Cesari



These are the first engineering data cubes for each of the twelve MRS spectral bands, illustrating the astrometric registration and image quality for observations of HD 37122. In each panel the dashed cyan circle shows a 1 arcsecond radius region around the expected location of the star in celestial coordinates. While the star is bright at short wavelengths it fades toward longer wavelengths, where the MRS also detects thermal emission from Webb's primary mirror. Credit: NASA, ESA, and the MIRI Consortium


NASA's James Webb Space Telescope team continues to work its way through the 17 science instrument modes. This week they checked off numbers (5) NIRCам grism time series and (4) imaging time series, both used to study exoplanets and other time-variable sources; (12) NIRISS aperture masking interferometry mode, for direct detection of a faint object that is very close to a bright one; (11) NIRISS wide-field slitless

spectroscopy, for studying distant galaxies; and (9) NIRSpec bright-object time series, for studying exoplanets. That totals seven modes approved to date, with 10 still to go.

This week, we are featuring MIRI's medium-resolution spectroscopy mode and sharing our first spectroscopic engineering data. We asked two of the MIRI commissioning team members—David Law, of the Space Telescope Science Institute (STScI), and Alvaro Labiano, of the Centro de Astrobiología (CAB)—to explain this mode to us:

"One of Webb's most complex instrument modes is with the MIRI Medium Resolution Spectrometer (MRS). The MRS is an integral-field spectrograph, which provides spectral and [spatial information](#) simultaneously for the entire field of view. The spectrograph provides three-dimensional 'data cubes' in which every pixel in an image contains a unique spectrum. Such spectrographs are extremely powerful tools to study the composition and kinematics of astronomical objects, as they combine the benefits of both traditional imaging and spectroscopy.

"The MRS is designed to have a spectral resolving power (observed wavelength divided by the smallest detectable wavelength difference) of about 3,000. That is high enough to resolve key atomic and molecular features in a variety of environments. At the highest redshifts, the MRS will be able to study hydrogen emission from the first galaxies. At lower redshifts, it will probe molecular hydrocarbon features in dusty nearby galaxies and detect the bright spectral fingerprints of elements such as oxygen, argon, and neon that can tell us about the properties of ionized gas in the interstellar medium. Closer to home, the MRS will produce maps of spectral features due to water ice and simple organic molecules in [giant planets](#) in our own solar system and in planet-forming disks around other stars.



This portion of the MIRI MRS wavelength range shows engineering calibration data obtained of the Seyfert galaxy NGC 6552 (red line) in the constellation Draco. The strong emission feature is due to molecular hydrogen, with an additional weaker feature nearby. The blue line shows a lower spectral resolution Spitzer IRS spectrum of a similar galaxy for comparison. The Webb test observations were obtained to establish the wavelength calibration of the spectrograph. Credit: NASA, ESA, and the MIRI Consortium

"In order to cover the wide 5 to 28 [micron](#) wavelength range as efficiently as possible, the MRS integral field units are broken up into twelve individual wavelength bands, each of which must be calibrated individually. Over the past few weeks, the MIRI team (a large international group of astronomers from the U.S. and Europe) has been focusing primarily on calibrating the imaging components of the MRS.

They want to ensure that all twelve bands are spatially well aligned with each other and with the MIRI Imager, so that it can be used to place targets accurately into the smaller MRS field of view. We show some early test results from this alignment process, illustrating the image quality achieved in each of the twelve bands using observations of the bright K giant star HD 37122 (located in the southern sky near the Large Magellanic Cloud).

"Once the spatial alignment and image quality of the several bands are well characterized, the MIRI team will prioritize calibrating the spectroscopic response of the instrument. This step will include determining the wavelength solution and spectral resolution throughout each of the twelve fields of view using observations of compact emission-line objects and diffuse planetary nebulae ejected by dying stars. We show the exceptional spectral resolving power of the MRS with a small segment of a spectrum obtained from recent engineering observations of the active galactic nucleus at the core of Seyfert galaxy NGC 6552. Once these basic instrument characteristics are established, it will be possible to calibrate MRS so that it is ready to support the wealth of Cycle 1 science programs due to start in a few short weeks."

Provided by NASA

Citation: Webb's mid-infrared spectroscopy will reveal molecules, elements (2022, June 17) retrieved 26 April 2024 from

<https://phys.org/news/2022-06-webb-mid-infrared-spectroscopy-reveal-molecules.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.