

Fossils confirm early diversification of spiny plants in central Tibet

July 5 2022, by Zhang Nannan



A typical spiny plant Gleditsia microphylla (Fabaceae) native in Asia. Credit: ZHANG Xinwen

Spinescence (a general term for the phenomena of spines, prickles, and thorns on plants) is an important functional trait shared by numerous plant families worldwide and mainly provides physical protection against vertebrate herbivores. Even though spiny plants are distributed worldwide, our understanding of their evolutionary history remains



incomplete, largely due to a dearth of fossil records.

In a study published in *Nature Communications*, researchers from the Xishuangbanna Tropical Botanical Garden (XTBG), the Institute of Vertebrate Paleontology and Paleoanthropology of the Chinese Academy of Sciences (CAS), the University of Bristol, and the Open University of the United Kingdom have reported exceptionally rich assemblages of plant spine fossils collected from late Eocene (about 39 million years ago) sediments in central Tibet.

These fossils confirm an early diversification of spiny <u>plants</u> in the Tibetan region contemporaneous with the emergence of open, semiwooded habitats by the late Eocene and early in the transition of central Tibet to full plateau formation.

The researchers documented a total of 44 spine-bearing fossil specimens collected from two fossil localities within the central Tibetan Bangong-Nujiang Suture Zone: the Dayu ($32^{\circ} 20'$ N, $89^{\circ} 46'$ E) and Xiede ($31^{\circ} 58'$ N, $88^{\circ} 25'$ E) localities. They classified fossil spines into a total of seven morphotypes comprising prickles and thorns according to two criteria: prickles originate from the epidermis of plant organs such as stems, leaves and petioles, whereas thorns are modified branches and contain internal vascular bundles.





Morphotypes of spiny fossils from the Eocene sediments of the central Tibetan Plateau. Credit: ZHANG Xinwen





A spiny fossil from the Eocene sediments of the central Tibetan Plateau. Credit: ZHANG Xinwen

To investigate the emergence and early diversification of spiny plants, the researchers not only conducted molecular phylogenetic analyses, but also used proxy and modeling data to reconstruct the vegetation, climate, and herbivory that favored spiny plant evolution in central Tibet during the Eocene. In addition, they reconstructed the evolutionary history of spines across species of woody eudicots in Eurasia within the megaphylogeny of plants.

According to Zhang Xinwen of XTBG, the modeling and proxy data point to "a drying and cooling climate" in Tibet's central valley by the mid-Eocene. This change was accompanied by an increase in elevation



of about one kilometer about 47 million years ago.

Against this backdrop, the plant fossils record the beginnings of vegetation opening-up under a drying/cooling climate as the modern Tibetan Plateau grew from the late Eocene onwards. The observed spinescence marks the early development of physical defense mechanisms against large herbivore feeding pressure in the region.

"Our study shows that regional aridification and expansion of herbivorous mammals may have driven the diversification of functional spinescence in central Tibetan woodlands, about 24 million years earlier than similar transformations in Africa," said Su Tao of XTBG.

More information: Xinwen Zhang et al, Rapid Eocene diversification of spiny plants in subtropical woodlands of central Tibet, *Nature Communications* (2022). DOI: 10.1038/s41467-022-31512-z

Provided by Chinese Academy of Sciences

Citation: Fossils confirm early diversification of spiny plants in central Tibet (2022, July 5) retrieved 17 July 2024 from <u>https://phys.org/news/2022-07-fossils-early-diversification-spiny-central.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.