


New model developed to predict landslides along wildfire burn scars

July 27 2022



This illustration shows the steps leading to a debris flow: drought, followed by wildfires, followed by intense precipitation can lead to a fast-moving landslide with the ability to wipe out roads and homes. Credit: Tierney Acott/Institute of Sustainability and Energy at Northwestern

A wildfire followed by an intense rainstorm is often a recipe for disaster. Without vegetation to cushion rainfall, water runoff can turn into a fast-moving, highly destructive landslide, called a "debris flow," which often has the power to wipe out cars, homes and highways—sometimes resulting in casualties.

Northwestern University researchers have augmented a physics-based [numerical model](#) to investigate and predict areas susceptible to [debris](#) flows. This augmented model eventually could be used in an early warning system for people living in high-risk areas, enabling them to evacuate before it's too late. Information from [model simulations](#) also could be used to design new infrastructure—such as diversion bars that deflect fast-moving water away from homes and roads—for high hazard zones.

The research will be published on July 27 in the journal *Natural Hazards and Earth System Sciences*.

"People want to know about their immediate and future risk," said Northwestern's Daniel Horton, the study's senior author. "Although it's not yet to operational standards, this modeling framework could one day be instrumental in forecasting where debris flows are likely to occur and deciding who needs to be evacuated."

Horton is an assistant professor of Earth and planetary sciences in Northwestern's Weinberg College of Arts and Sciences, where he also leads the Climate Change Research Group. Chuxuan Li, a Ph.D. candidate in Horton's laboratory, is the paper's first author.

Increasing debris flow 'ingredients'

As climate change increases the likelihood of wildfires and [heavy rains](#), debris flows also might become more common. [Just last year](#), a debris

flow tore out a 150-foot section of roadway along Highway 1 near Big Sur, California. The picturesque two-lane highway was closed for three months, cutting off tourism to the region.

The disaster resulted from a one-two punch: A wildfire that burned from August through December 2020, followed by an atmospheric river—a long, thin string of concentrated moisture in the sky—that dumped nearly a foot of rain the following January.

Following intense rainfall, areas with wildfire burn scars are more susceptible to flash flooding. In a typical situation, vegetation interacts with rain, which then soaks into the ground. But if an area is burned, vegetation is gone, so it can no longer intercept the sudden influx of moisture. Even worse: In some regions, waxes from burned vegetation melts, enters the soil and then solidifies. These waxes create a barrier for rain, blocking it from penetrating the soil. Instead, rain runs over the top of the soil, picking up rocks, mud, trees and even cars to become a debris flow.

"Because of human-caused climate change, the ingredients that lead to debris flows are certainly becoming more common," Horton said. "Due to intensified precipitation, drought and wildfires, debris flow conditions are particularly prevalent in California."

Pinpointing susceptible areas

To develop the new model, Horton's team adapted the National Oceanic Atmospheric Administration's National Water Model, which is used to simulate streamflow and [water resources](#) across the continental United States.

"The National Water Model is a community, open-source model, so researchers use it to probe all sorts of pertinent water resource and

hazard questions," Horton said. "But one thing the model didn't explicitly output was overland flow—water that runs across the land after rainfall. We modified the model to look at how extreme precipitation moves over land surfaces, which allows us to investigate how susceptible certain land surfaces, and in particular burn scars, are to debris flow."

By adjusting vegetation type (dense vs. barren, for example) and soil properties, the researchers were able to replicate how water moves over wildfire burn scars. The team then validated the model-simulated hydrological conditions using soil moisture and streamflow observations.

The resulting simulation found that burned areas cause [water](#) to flow much faster and in greater volumes compared to unburned land. When simulating the debris flow that took a bite out of Highway 1, for example, the researchers found overland flow volume increased eight-fold and peak flow tripled compared to land without burn scar characteristics.

"An advantage of our model is that it uses precipitation as it would fall in the real world," Horton said. "We're simulating spatially complex and time-evolving precipitation and how that precipitation interacts with the land surface."

"Our physics-based hydrological [model](#) opens up new possibilities for better understanding of burn scar hydrology and debris flow susceptibility," Li said. "Given the increasing likelihood of debris flow conditions due to [climate change](#), developing models to better understand past events and predict future events is essential."

The study is titled "Augmentation of WRF-Hydro to simulate overland flow- and streamflow-generated debris [flow](#) susceptibility in burn scars."

More information: Chuxuan Li et al, Augmentation and Use of WRF-

Hydro to Simulate Overland Flow- and Streamflow-Generated Debris Flow Hazards in Burn Scars, *Natural Hazards and Earth System Sciences* (2021). [DOI: 10.5194/nhess-2021-345](https://doi.org/10.5194/nhess-2021-345)

Provided by Northwestern University

Citation: New model developed to predict landslides along wildfire burn scars (2022, July 27) retrieved 13 March 2024 from <https://phys.org/news/2022-07-landslides-wildfire-scars.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.