

New study challenges old views on what's 'primitive' in mammalian reproduction

July 25 2022, by James Urton



An artistic rendering of multituberculates from the genus *Mesodma* — a mother with her litter of offspring — who lived in western North America about 60 to 70 million years ago. Fossil evidence indicates that these creatures were the most abundant mammals in western North America just before and directly after the mass extinction event 66 million years ago that killed off the dinosaurs. Credit: Andrey Atuchin

It's hard to imagine life on Earth without mammals. They swim in the depths of the ocean, hop across deserts in Australia and travel to the moon.

This diversity can be deceiving, at least when it comes to how mammals create the next generation. Based on how they reproduce, nearly all mammals alive today fall into one of two categories: placental mammals and marsupials. Placentals, including humans, whales and rodents, have long gestation periods. They give birth to well-developed young—with all major organs and structures in place—and have relatively short weaning periods, or lactation periods, during which young are nursed on milk from their mothers. Marsupials, like kangaroos and opossums, are the opposite: They have short gestation periods—giving birth to young that are little more than fetuses—and long lactation periods during which offspring spend weeks or months nursing and growing within the mother's pouch, or marsupium.

For decades, biologists saw the marsupial way of reproduction as the more "primitive" state, and assumed that placentals had evolved their more "advanced" method after these two groups diverged from one another. But new research is testing that view. In a paper published July 18 in *The American Naturalist*, a team led by researchers at the University of Washington and its Burke Museum of Natural History and Culture present evidence that another group of mammals—the extinct

multituberculates—likely reproduced in a placental-like manner. Since multituberculates split off from the rest of the mammalian lineage before placentals and marsupials evolved, these findings question the view that marsupials were "less advanced" than their placental cousins.

"This study challenges the prevalent idea that the placental reproductive strategy is 'advanced' relative to a more 'primitive' marsupial strategy," said lead author Lucas Weaver, a postdoctoral researcher at the University of Michigan who conducted this study as a UW doctoral student. "Our findings suggest that placental-like reproduction either is the ancestral reproductive route for all mammals that give birth to live young, or that placental-like reproduction evolved independently in both multituberculates and placentals."

Multituberculates arose about 170 million years ago in the Jurassic. Most were small-bodied creatures, resembling rodents. For much of their history, multituberculates were the most abundant and diverse group of mammals. But scientists know very little about their [life history](#), including how they reproduced, because of their generally poor fossil record. The last multituberculates died out about 35 million years ago.


Weaver reasoned that the microscopic structure of fossilized bone tissues can house useful life-history information about multituberculates, such as their growth rate. Working under co-author Gregory Wilson Mantilla, a UW professor of biology and curator of vertebrate paleontology at the Burke Museum, Weaver and his colleagues obtained cross sections of 18 fossilized femurs—the [thigh bone](#)—from multituberculates that lived approximately 66 million years ago in Montana.

All 18 samples showed the same structural organization: a layer of disorganized bone "sandwiched" between an inner and outer layer of organized bone. Disorganized bone, or woven bone, indicates rapid


growth and is so named because, under a microscope, the layers of bone tissue are laid out in a crisscrossed fashion. In organized bone, which reflects slower growth, layers are parallel to one another.

The researchers then examined femoral cross sections taken from 35 small-bodied mammalian species that are living today—28 placentals and seven marsupials, all from Burke Museum collections. Nearly all of the placental femurs showed the same "sandwich" organization as the multituberculates. But all of the marsupial femurs consisted almost entirely of organized bone, with only a sliver of disorganized bone.

Virginia opossum (marsupial)



Eastern chipmunk (placental)



Multituberculate



The three images are cross sections of femurs from a marsupial (the Virginia opossum, left), a placental (the eastern chipmunk, center) and a 66-million-year-old multituberculate fossil (right). The opossum femur has a thick layer of organized bone in the outermost cortex (labeled "POB" for periosteal organized bone), with little disorganized bone (labeled "DB"). In the chipmunk and multituberculate femurs, a layer of disorganized bone ("DB") is "sandwiched" between layers of organized bone ("POB" and "EOB," which stands for endosteal organized bone). Scale bar is 0.1 millimeters. The multituberculate specimen (UWBM 70536) is likely a member of the genus *Mesodma*. Credit: Henry Fulghum/Lucas Weaver/University of Washington

The team believes that this stark difference likely reflects their divergent life histories.

"The amount of organized bone in the outermost layer, or cortex, of the femur strongly correlates with the length of the lactation period," said Weaver. "Marsupials have long lactation periods and a lot of organized bone in the outermost cortex. The opposite is true for placentals: a short lactation period and much less organized bone in the outermost cortex."

The outermost layer of organized bone was laid down after birth as the femur's diameter increased. For tiny [marsupial](#) newborns, bones must grow much more to reach adult size, so they deposit a greater amount of outer organized [bone](#) compared to placentals, according to Weaver.

"This is compelling evidence that multituberculates had a long gestation and a short lactation period similar to [placental mammals](#), but very different from marsupials," said Weaver.

Based on this correlation, the researchers estimate that multituberculates had a lactation period of approximately 30 days—similar to today's rodents.

These findings cast further doubt on an old view that marsupials have a "more primitive" and placentals a "more advanced" reproductive strategy. The common ancestor of multituberculates, placentals and [marsupials](#) may have had a placental-like mode of reproduction that was retained by placentals and multituberculates. Alternatively, multituberculates and placentals could have evolved their long-gestation and short-lactation reproductive methods independently.

Future studies of multituberculate life history may clarify which explanation is true, as well as other outstanding questions of this, and other, ancient branches of our mammalian family tree.

"The real revelation here is that we can cut open fossil bones and examine their microscopic structures to reconstruct the intimate life history details of long-extinct mammals," said Wilson Mantilla. "That's really incredible to me."

More information: Lucas N. Weaver et al, Multituberculate Mammals Show Evidence of a Life History Strategy Similar to That of Placentals, Not Marsupials, *The American Naturalist* (2022). [DOI: 10.1086/720410](https://doi.org/10.1086/720410)

Provided by University of Washington

Citation: New study challenges old views on what's 'primitive' in mammalian reproduction (2022, July 25) retrieved 27 April 2024 from <https://phys.org/news/2022-07-views-primitive-mammalian-reproduction.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--