


Organic matter accumulation in oxygenated lakes

August 25 2022



Conceptual figure showing differences between open (A) and closed (B) basin lakes and the production of recalcitrant DOM (RDOM). Although based on work in the present study, this conceptual model likely applies to most freshwater systems with water residence time being an important variable. Open basins are strongly connected to their watershed and have short residence times which provides CDOM and promotes microbial diversity. CDOM absorbs solar radiation which decreases solar radiation to increased depths in the water column. CDOM absorption also leads to the production of reactive oxygen

species which facilitate DOC degradation and consumption by bacteria. Most of the DOC production by phytoplankton and macrophytes is labile (LDOM) and degraded by photo-chemical and microbial processes into primarily other labile compounds and CO₂, leading to little net ecosystem production due to a balance between photosynthesis (P) and respiration (R). In closed basin, long residence-time lakes, connections with the watershed are cut off with little CDOM and microbial inputs. Therefore, light levels are higher in the lake, increasing the role of photosynthetic processes while decreasing the diversity of microbes capable of degrading DOC, enabling $P > R$. These conditions coupled with longer time scales for DOM degradation lead to increased pool sizes of RDOM. Credit: *Limnology and Oceanography Letters* (2022). DOI: 10.1002/lol2.10265

When we burn fossil fuels, it not only produces carbon dioxide, a driver of climate change, but it also consumes the oxygen we breathe. However, the amount of oxygen in our atmosphere produced by plants is nearly balanced by the amount consumed by animals, keeping it at around 21% of the atmosphere. This raises a big question relevant to our survival and the future of biodiversity: what keeps the levels of oxygen in our atmosphere relatively constant?

Oxygen helps break down [organic matter](#) to release [carbon dioxide](#)—a process you can see in a backyard compost pile. Yet in some places on Earth, organic matter such as plant debris can persist for thousands of years despite the presence of abundant oxygen. College of Biological Sciences professor James Cotner wants to better understand why this occurs and its implications for [carbon sequestration](#) and climate change.

In a recent study published in the journal *Limnology and Oceanography Letters*, Dr. Cotner and co-authors N.J. Anderson and Christopher Osburn sampled lakes in Greenland where dissolved [organic compounds](#) can accumulate to concentrations that are 200 times greater than concentrations in the oceans. Some of these lakes sit right next to lakes

with much lower concentrations, and they wanted to understand why organic matter is preserved in some lakes but not others. They measured the salinity of the lakes to determine how connected the lakes were to their watersheds and used radiocarbon dating to measure the age of the organic matter. "Our work seems to suggest that hydrology and sunlight can have a big effect on sequestration." says Cotner.

The researchers found that:

- Less salty lakes that were well-connected to surrounding watersheds degraded organic matter much more quickly than lakes that were somewhat isolated, i.e., more salty.
- Exposure to sunlight and the production of metabolites by microbes when they are starved for nutrients likely facilitate organic matter accumulation in the less-connected lakes.
- Hydrology is a key part of the carbon cycle on Earth's surface, and both natural and human-driven processes like agriculture impact hydrologic connectivity with implications for both the [carbon cycle](#) and oxygen in our atmosphere.
- "We were somewhat shocked to find that the age of the dissolved organic matter correlated extremely well with the salinity of the [lake](#) water, suggesting that very old organic matter can persist in lakes that are not well-connected to their surrounding landscapes." said John Anderson, a co-author from Loughborough University.

Further research in this area could reveal more about how carbon sequestration occurs in nature, which could have implications for human efforts at carbon sequestration as well. "Our future work will be focusing on the importance of tannins, humic compounds and nutrients as well as the role of different soil microbes to the degradation of organic matter in freshwater," says Cotner.

More information: James B. Cotner et al, Accumulation of recalcitrant dissolved organic matter in aerobic aquatic systems, *Limnology and Oceanography Letters* (2022). [DOI: 10.1002/lol2.10265](https://doi.org/10.1002/lol2.10265)

Provided by University of Minnesota

Citation: Organic matter accumulation in oxygenated lakes (2022, August 25) retrieved 13 April 2024 from <https://phys.org/news/2022-08-accumulation-oxygenated-lakes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.