

# Climate change leads to invasive insect expansion on US West Coast

August 10 2022

---



A collage of oak galls created by oak gall wasps. Credit: Kirsten Prior

Climate change has led to warming temperatures in the Pacific

Northwest, leading some insect species to expand their range into more northerly oak savannas, according to new research from Binghamton University, State University of New York.

Side by side, Dylan Jones displayed photos of two oak leaves. One was healthy and green, dotted by the occasional gall, a structure made by a herbivorous species of oak gall wasp. The other leaf was yellowed and tattered, the victim of an insect population without predatory checks and balances. Climate change has led to warming temperatures in the Pacific Northwest, leading species such as *Neuroterus saltatorius* to expand their range into more northerly oak savannas.

"In the native range, you might find a handful of galls on a single leaf. In the expanded range, sometimes you're finding thousands on a single tree," said Assistant Professor of Biological Sciences Kirsten Prior. "This is pretty prevalent throughout Vancouver Island."

Jones, a Binghamton University doctoral candidate in biological sciences and Clifford D. Clark Diversity Fellow, is the lead author of a research paper recently published in the *Journal of Animal Ecology* on the situation. Co-authors on "Latitudinal gradient in species diversity provides high niche opportunities for a range-expanding phytophagous insect" include Prior, field technician Julia Kobelt, then-undergraduate Jenna Ross and Assistant Professor of Biological Sciences Thomas Powell.

Oak savannas are grassy and shrubby areas that feature oaks as the dominant tree species. The oak species in question—*Quercus garryana*—requires a dry environment. As a result, oak savannas are often found in the rain shadow of the West Coast's mountain range, Prior explained.

A diverse and ecologically important group of [trees](#) throughout North

America, oaks host a wide diversity of insect species, such as oak gall wasps. These wasp species form growths known as galls that can be striking in their varied forms, from those that resemble large apples to others with colorful spikes reminiscent of sea urchins.

Interestingly, Alfred Kinsey—yes, that Kinsey—studied oak gall wasps before transitioning his field of study to human sexuality.

"There has been a long fascination by biologists and amateurs for this group of species because they're pretty charismatic," Prior said. "You can go to an oak and see all these structures on it."

In addition to their herbivorous creators, these [galls](#) host other species of insects, including a dazzling variety of [parasitoid wasps](#), one of the most diverse and ecologically important groups in the animal kingdom, the latter for their role in keeping insect pests in check. Parasitoid wasps lay their eggs in the gall; after hatching, their [larvae](#) then eat the oak gall wasp larvae.

Oak gall wasps are highly diverse in North America, but not well-documented; their evolutionary relationships and even the identification of some species remain unknown. A consortium of scientists across the continent is working to change that; Jones and Prior are part of this effort.

"It's important to continue to document biodiversity. We still haven't described a good portion of the biodiversity on Earth, particularly with insects," Prior said.

## **Ecosystem invaders**

The researchers checked their study sites on three different occasions during the course of that summer. Some were quite remote, involving

trips down dirt logging roads, or located on Bureau of Land Management property or on preserves associated with Indigenous communities. Others were suburban, located a short jaunt from cities.

Because of urbanization, few oak savannas remain on Vancouver Island; those that are left are highly documented and tended by landowners.

"We've had long relationships with lots of landowners there that allow us to work on their property," Prior said. "Some of them are so jazzed to have researchers there."

Biodiversity tends to work on a latitudinal gradient, Jones added: The closer you are to the equator, the more species you have. A similar situation holds true when it comes to higher altitudes. When one species can expand its range due to warming temperatures, it can move into areas without a diversity of predators and competitors, eventually overwhelming the ecosystem.

The case of the oak gall [wasps](#) highlights the importance of biodiversity and the potential long-term ramifications of climate change, the researchers point out.

"Biodiversity can be really important in potentially protecting areas from invading species," Jones said. "If we have strong competitors and predators, this might make areas less susceptible to invading species."

**More information:** Latitudinal gradient in species diversity provides high niche opportunities for a range-expanding phytophagous insect, *Journal of Animal Ecology* (2022). [DOI: 10.1111/1365-2656.13780](https://doi.org/10.1111/1365-2656.13780)

Provided by Binghamton University

Citation: Climate change leads to invasive insect expansion on US West Coast (2022, August 10)  
retrieved 2 May 2024 from

<https://phys.org/news/2022-08-climate-invasive-insect-expansion-west.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.