


How life on land learned to breathe

August 12 2022



(a) Partial skeletal reconstruction of *Gorgonops torvus* in lateral aspect and (b) schematic reconstruction of the pectoral girdle, sternum, interclavicle and humerus with hypothetical position of *m. pectoralis* in ventral aspect. (c) Photograph and (d) interpretative drawing of *Gorgonops torvus* (SAM-PK-K10591), highlighting the pectoral region in an orange rectangle as an (e)

photograph and (f) interpretative drawing (e and f after further preparation to reveal the interclavicle). For explanation of the taphonomic disturbance of the specimen, see Materials and methods section. Abbreviations: cl, clavicle; cr, cervical ribs; cv, cervical vertebrae; fe-L, left femur; fi-L, left fibula; h, partial humeri; icl, interclavicle; m-L, left manus; 'ma', 'manubrium'; p-L, left pes; pc-R, right pectoral girdle; pel?, partial pelvic girdle?; pel-L, left pelvic girdle; r-L, left radius; r-R, right radius; ri, ribs; sc-L; left scapula; sc-R, right scapula; st1-3, sternebra 1–3; ti-L, left tibia; u-L, left ulna; v, vertebrae. Credit: *Scientific Reports* (2022). DOI: 10.1038/s41598-022-17492-6

Scientists from the Museum für Naturkunde Berlin, together with collaborators from South Africa and the U.S., have revealed a new chapter in the history of the mammalian breastbone. Their study of a 260-million-year-old fossil shows that its breastbone was divided into a series of segments, like that of modern mammals. This discovery indicates that prerequisites of the mammalian way of walking and breathing were already present in some distantly related Permian ancestors.

In a recent study, led by Ph.D. candidate Eva-Maria Bendel of the Museum für Naturkunde Berlin and the Humboldt-University Berlin, the evolutionary history of the mammalian [breastbone](#) has been traced. Together with experts from the North Carolina Museum of Natural Sciences, the University of Chicago, and the Iziko South African Museum, the team links the anatomy of a group of mammalian forerunners with the way of life of today's mammals.

Based on an especially complete specimen at the Iziko [museum](#) in Cape Town, South Africa, Bendel identified the oldest recorded sternum consisting of multiple elements, just like in modern mammals. "I noticed right away that the configuration in this gorgonopsian is something exceptional," she notes.

In the open access journal *Scientific Reports*, the team now describes the thorax and shoulder girdle of the Permian top predator *Gorgonops torvus*. The findings suggest one of two possibilities: (1) that the "[mammal](#)-like" condition arose first in the group Gorgonopsia but then was temporarily lost again on the mammalian stem; or (2) that this trait evolved independently in the two groups.

The presence of a segmental breastbone furthermore helps mammals to a more upright posture and erect gait, drastically enhancing their mobility. The researchers assume that the characteristic way mammals breathe by means of a diaphragm has already developed in their early ancestors. This may have aided gorgonopsians in becoming the dominant group of predators at top of the food chain in the late Permian.

More information: Eva-Maria Bendel et al, The earliest segmental sternum in a Permian synapsid and its implications for the evolution of mammalian locomotion and ventilation, *Scientific Reports* (2022). [DOI: 10.1038/s41598-022-17492-6](https://doi.org/10.1038/s41598-022-17492-6)

Provided by Museum für Naturkunde Berlin

Citation: How life on land learned to breathe (2022, August 12) retrieved 11 May 2024 from <https://phys.org/news/2022-08-life.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.