

# Lime shows promise for controlling clubroot in canola crops

August 8 2022, by Bev Betkowski

---



Nicole Fox found that treating soil with lime may offer an additional line of defence to help protect canola crops against clubroot, along with use of plants

bred for genetic resistance to the disease. Credit: University of Alberta

Lime is showing promise as an additional way to help manage clubroot, a deadly disease in canola crops, University of Alberta research shows.

Spot-treating soil with the mineral reduced the overall occurrence and severity of the disease by 35 to 91%, growth experiments showed.

The finding, published in the *Canadian Journal of Plant Pathology*, could give farmers an important option for managing clubroot in their fields, alongside current use of [canola](#) plants bred to resist the disease.

Lime has traditionally been used to manage clubroot in related plants such as cabbages for market gardens, but not on a large scale in canola crops.

"Genetics are our first line of defense, but plant resistance can erode or break down, so we need to find every possible option to help control the clubroot pathogen in what is an important cash crop for Canada," says researcher Nicole Fox.

As a non-genetic management practice, liming treatments could help combat all strains of clubroot in canola, adds Fox, who conducted the study to earn a master of science in plant biosystems from the Faculty of Agricultural, Life & Environmental Sciences.

Spot treatments could help control contaminated areas of a field or stem the spread of clubroot into a new field.



Side-by-side comparison of clubroot-infested canola plants growing in soil treated with lime (right) and plants growing in untreated soil (left). Credit: University of Alberta

The lime works by neutralizing the highly [acidic soil](#) preferred by clubroot, reducing the likelihood of spore germination and plant infection.

One of the first studies to test hydrated lime in the field in Canada, the research also showed that the product was more effective at managing clubroot in canola than granulated limestone, another form of the mineral that is more commonly used to treat agricultural soil.

Applying moderate to high amounts of the powdered lime resulted in

canola plants that were still productive even if they were already infected by clubroot. The plants also released fewer spores of the clubroot pathogen back into the soil.

Lowering the [acidity levels](#) also increases the soil's general health—an important benefit to liming, considering Alberta has about one million acres of strongly acidic and 4.5 million acres of moderately acidic cropland soils, Fox notes.

But hydrated lime's effectiveness does hinge on certain factors, like the interval between application and seeding, so it needs fine-tuning before it could become a practical tool, says U of A plant pathologist and study co-author Stephen Strelkov, who supervised the research.

"While lime showed good potential for clubroot management, the results varied. Sometimes the treatments provided very good control; other times they didn't. So we need further research to work out some details."

Canola bred to be genetically resistant is still the most effective tool against clubroot, he adds, but options like using hydrated lime could help improve the "durability of resistance and overall sustainability" of disease management.

"There may be situations, if the efficacy of [lime](#) is consistent and the costs of application reasonable, that it could be used on a larger scale."

**More information:** Nicole M. Fox et al, Evaluation of lime products for clubroot (*Plasmodiophora brassicae*) management in canola (*Brassica napus*) cropping systems, *Canadian Journal of Plant Pathology* (2021). [DOI: 10.1080/07060661.2021.1940590](https://doi.org/10.1080/07060661.2021.1940590)

Provided by University of Alberta

Citation: Lime shows promise for controlling clubroot in canola crops (2022, August 8) retrieved 26 April 2024 from <https://phys.org/news/2022-08-lime-clubroot-canola-crops.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.