


Keeping toxic cadmium out of rice, the genetic way

August 30 2022



In Pokkali, due to duplicated OsNramp5, the roots take up more Mn and Cd into the root cells compared with other rice varieties (for example, Koshihikari). Most Cd taken up through OsNramp5 is sequestered into vacuoles by OsHMA3, while most Mn is translocated to the shoots through OsMTP9. Higher Mn concentration in the root cells competes with Cd for an unidentified efflux transporter, resulting in decreased loading of Cd to the xylem and subsequently

to the shoots and grains. The dotted arrow shows decreased Cd loading by high Mn. The cyan and purple colors indicate Mn and Cd, respectively. Credit: Jian Feng Ma, Okayama University

Rice is a staple food for nearly half the world's population. However, it accumulates more cadmium from the soil than other cereals like barley and wheat. Reports estimate that 40–65% of our total intake of cadmium, a toxic heavy metal, is from rice. Eating cadmium-contaminated rice poses a serious health risk to humans, with conditions like Itai-itai disease being associated with high cadmium intake.

Efforts have been previously made to reduce the quantity of [cadmium](#) in rice through methods like importing clean soil, [water management](#), and mixing contaminated soil with biochar and lime. However, these methods are time-consuming and expensive. To remedy this, scientists have turned to cross-breeding to cultivate rice that accumulate less cadmium.

"We have been working on the mechanisms of cadmium accumulation in rice and barley for a long time and have identified several key genes involved in its accumulation," says Prof. Jian Feng Ma, who is affiliated to the Institute of Plant Sciences and Resources at Okayama University, Japan. Recently, Prof Ma published a paper in the journal *Nature Food* detailing the genetic mechanisms that play a role in this process.

After examining 132 accessions of rice, Prof Ma and the members of his research group found that the gene, OsNramp5, when duplicated in tandem, reduced the accumulation of cadmium in Pokkali, a variety of rice that has been cultivated for 3000 years in Kerala, India. According to previous reports, OsNramp5 encodes a cadmium and manganese transporter protein in rice. The same gene, when duplicated in tandem,

turns to increase the uptake of both the minerals into the root cells. Consequently, manganese competes with cadmium in the cells for translocation to the shoots, which in turn reduces the accumulation of cadmium in these parts.

The scientists found that out of the 132 accessions of rice, the tandem duplication of OsNramp5 was naturally found only in Pokkali, which can grow in salt-laden coastal soil.

The researchers also noted that the spatial expression level of OsNramp5 was always around two-fold higher in the roots of Pokkali than that in the roots of Koshihikari.

As Pokkali stores extremely low cadmium in its shoots, the scientists introgressed (a term for the transfer of genetic information across species) the duplicated OsNramp5 gene in Koshihikari, a variety of rice that is very popular in Japan but accumulates relatively high levels of cadmium. Explaining how targeted breeding can help humans, Prof. Ma says that they "identified a gene responsible for differential accumulation of cadmium in rice grain based on natural variations in cadmium accumulation. Then, we applied this gene to successfully breed rice cultivars with low cadmium accumulation in the grain."

The team found that the Koshihikari cultivar with the duplicated gene accumulated significantly lower amounts of cadmium without comprising on the grain quality or yield.

Recounting the benefits of a low cadmium-accumulating rice variety, Prof. Ma explains that "cadmium is a toxic heavy metal and threatens our health through the food chain. Our study provided a useful material for breeding varieties of rice with low cadmium accumulation, which contributes to produce safe and healthy food. We hope that this gene will be widely used in breeding different [rice](#) cultivars with low cadmium

accumulation. This will protect us from cadmium poisoning."

More information: Jian Ma, Duplication of a manganese/cadmium transporter gene reduces cadmium accumulation in rice grain, *Nature Food* (2022). DOI: [10.1038/s43016-022-00569-w](https://doi.org/10.1038/s43016-022-00569-w).
www.nature.com/articles/s43016-022-00569-w

Provided by Okayama University

Citation: Keeping toxic cadmium out of rice, the genetic way (2022, August 30) retrieved 23 May 2024 from <https://phys.org/news/2022-08-toxic-cadmium-rice-genetic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.