

## **Study tracks plant pathogens in leafhoppers from natural areas**

August 2 2022



Leafhoppers that are known—or are likely—to transmit phytoplasmas to plants include, clockwise, from top left, species of the genera Hishimonoides, Macrosteles, Amplicephalus, Osbornellus and Amplicephalus. The leafhopper on the lower right, Osbornellus auronitens, was found for the first time to harbor a phytoplasma strain. Credit: Christopher Dietrich

Phytoplasmas are bacteria that can invade the vascular tissues of plants, causing many different crop diseases. While most studies of



phytoplasmas begin by examining plants showing disease symptoms, a new analysis focuses on the tiny insects that carry the infectious bacteria from plant to plant. By extracting and testing DNA from archival leafhopper specimens collected in natural areas, the study identified new phytoplasma strains and found new associations between leafhoppers and phytoplasmas known to harm crop plants.

Reported in the journal *Biology*, the study is the first to look for phytoplasmas in insects from <u>natural areas</u>, said Illinois Natural History Survey postdoctoral researcher Valeria Trivellone, who led the research with INHS State Entomologist Christopher Dietrich. It also is the first to use a variety of molecular approaches to detect and identify phytoplasmas in leafhoppers.

"We compared traditional molecular techniques with next-generation sequencing approaches, and we found that the newer techniques outperformed the traditional ones," Trivellone said. These methods will allow researchers to target more regions of the phytoplasma genomes to get a clearer picture of the different bacterial strains and how they damage plants, she said.

"One thing that is really novel about this study is that we've focused on the vectors of <u>disease</u>, on the leafhoppers, and not on the plants," Dietrich said. The standard approach of looking for phytoplasmas in plants is much more labor-intensive, requiring that scientists extract the DNA from a plant that appears to be diseased and checking for phytoplasmas, he said.

"But even when you identify the phytoplasma, you don't know what leafhopper or other vector transmitted it to the plant," Dietrich said. "So researchers must go back out into the field to collect all potential insect vectors. Then they do transmission experiments, where they let the leafhoppers feed on an infected plant and then put them on an



uninfected plant to see if it catches the disease."



Researchers used both traditional and newer genetic sequencing techniques to look for phytoplasmas in leafhoppers. Credit: Fred Zwicky

Because this research is laborious and slow, "we still don't have a good idea of which insects are spreading most phytoplasmas between plants," Dietrich said. "That really limits your ability to set up an effective management strategy."

For the new study, the researchers turned to leafhopper specimens in the INHS insect collection. Dietrich had collected many of these insects over a period of 25 years as part of his work classifying their genetic



relatedness and evolution. The researchers examined 407 leafhopper species collected around the world in areas less disturbed by human development. The specimens came from North and South America, Africa, Europe, Asia and Australia.

The team extracted total DNA from the specimens and processed each one, using both traditional and newer sequencing approaches. The latter are less costly and more informative than traditional methods, the researchers report. Of the insects sampled, 41 tested positive for phytoplasmas, and the researchers obtained usable phytoplasma sequence data from 23 leafhoppers. The phytoplasmas included those that cause a disease known as aster yellows, which inhibits photosynthesis and reduces the productivity of several different <u>crop</u> <u>plants</u>. These phytoplasmas were found in several new species of leafhoppers never before identified as vectors of the disease.

"These leafhoppers may transmit the phytoplasmas to wild plants in natural areas," Trivellone said.

The study found phytoplasmas in regions of the world where such diseases had not been reported and identified several new strains of bacteria. It also found previously unreported associations between some phytoplasmas and species of leafhopper.





Dietrich examines leafhopper species collected over a period of 25 years and now in the INHS collection. Credit: Fred Zwicky

Scientists have no tools to target the bacteria in asymptomatic plants to prevent disease outbreaks, so controlling phytoplasmas involves the use of pesticides to kill the insect vectors.

"Because the insecticides are only partially specific to the target insects, they kill a variety of beneficial insects as well, which is not sustainable," Trivellone said.

"We're finding that there are lots of new phytoplasmas out there in nature that nobody's ever seen before," Dietrich said. "They don't cause <u>disease symptoms</u> in the native plants they've associated with for maybe



millions of years. They only start causing disease when they jump to a new host that has not been exposed to the phytoplasma before."

The new findings parallel those seen in emerging infectious diseases of humans originating in wildlife, Dietrich said. "This is why we need to look more broadly across nature and see what's out there."

**More information:** Valeria Trivellone et al, Comparison of Traditional and Next-Generation Approaches for Uncovering Phytoplasma Diversity, with Discovery of New Groups, Subgroups and Potential Vectors, *Biology* (2022). <u>DOI: 10.3390/biology11070977</u>

## Provided by University of Illinois at Urbana-Champaign

Citation: Study tracks plant pathogens in leafhoppers from natural areas (2022, August 2) retrieved 27 June 2024 from <u>https://phys.org/news/2022-08-tracks-pathogens-leafhoppers-natural-areas.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.