


Escaping atmosphere of hydrogen and helium from exoplanet studied with advanced simulations

September 19 2022, by Li Yuan



The stellar XUV and FUV/NUV SED, which is the spectrum of ϵ Eri taken from the MUSCLES Treasury Survey (France et al. 2016) and scaled at the orbital distance of WASP-52b. In panel (a), different solid lines represent the constructed spectra of different β_m . $\beta_m = 0.22$ corresponds to the reference spectrum, which is taken from the MUSCLES Treasury Survey. Credit: *The Astrophysical Journal* (2022). DOI: 10.3847/1538-4357/ac8793

Researchers from the Yunnan Observatories of the Chinese Academy of Sciences and their collaborators reproduced the observed transmission spectra of the exoplanet WASP-52b at different wavelength bands and

studied the properties of its hydrogen and helium atmosphere.

The results were published in *The Astrophysical Journal* on Sept. 13.

Close-in exoplanets receive intense high-energy radiation from their host stars, such as X-rays and extreme ultraviolet (XUV) radiation. In gas-rich exoplanets, the atmosphere may absorb this high-energy radiation, thus heating the atmosphere and causing it to expand to overcome the planet's gravitational potential and escape into the interstellar medium.

This phenomenon is known as planetary atmosphere escape, which can cause the loss of a large amount of material from the planet and has important effects on the composition, evolution and even the overall distribution of the planet.

The composition and properties of the planetary atmosphere can be studied by analyzing the absorption of spectral lines at different wavelength bands, for example, the optical band ($H\alpha$) and near-infrared band ($He\ \lambda 10830\text{\AA}$) lines, the so called [transmission](#) spectra.

In this study, the researchers used the hydrodynamic atmospheric escape model and the radiation transfer model to simulate the transmission spectra of WASP-52b. They introduced the Monte Carlo model to simulate the $Ly\alpha$ resonant scattering inside the exoplanetary atmosphere for the first time, by assuming that both the stellar $Ly\alpha$ radiation and the planetary atmosphere are in spherical.

Based on the distribution of the $Ly\alpha$ scattering rate $P\alpha$, the researchers calculated the $H\alpha$ absorption, which is caused by the [hydrogen atoms](#) in the first excited states. They also calculated the distribution of metastable helium atoms in detail and simulated the transmission [spectra](#) of hot Jupiter WASP-52b in the optical band ($H\alpha$) and near-infrared band ($He\ \lambda 10830\text{\AA}$).

They constrained the level of X-rays and extreme ultraviolet [radiation](#) received by the planet, as well as the hydrogen to helium abundance ratio in the planetary atmosphere, and revealed that [hydrogen](#) and helium originated from the escaping atmosphere. The findings can help to constrain the physical parameters of the [atmosphere](#) and to better understand its composition and structure.

More information: Dongdong Yan et al, Modeling the H α and He 10830 Transmission Spectrum of WASP-52b, *The Astrophysical Journal* (2022). [DOI: 10.3847/1538-4357/ac8793](https://doi.org/10.3847/1538-4357/ac8793)

Provided by Chinese Academy of Sciences

Citation: Escaping atmosphere of hydrogen and helium from exoplanet studied with advanced simulations (2022, September 19) retrieved 11 May 2024 from <https://phys.org/news/2022-09-atmosphere-hydrogen-helium-exoplanet-advanced.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--