


Researchers determine bright-magnetic relationship for decaying sunspots

September 26 2022, by Li Yuan



Comparison of the original HMI maps with deconvolved ones for the sunspot on NOAA AR 12662 observed on 2017 June 19 10:00 UT. The top row shows the difference of continuum intensity maps before and after applying the scattered-light correction. The white contours indicate the boundaries of the sunspot as seen in the original continuum intensity maps. The bottom row is similar, but for the vertical magnetic field (B_z) data. The third column shows the results of original data vs. the deconvolved data in the entire frames of the first and second columns. The blue circles mark the data of the sunspots. Credit: *The Astrophysical Journal* (2022). DOI: 10.3847/1538-4357/ac83b3

Prof. Yan Xiaoli from Yunnan Observatories of the Chinese Academy of Sciences and Li Qiaoling, a post-doctor from Yunnan University, have determined the bright-magnetic relationship for decaying sunspots for the first time.

Related results were published in *The Astrophysical Journal*.

The [brightness](#) of sunspots is an important parameter in the study of the total irradiance of the sun and a key index to verify the dynamics and magnetic properties of sunspots. The relationship between [sunspot](#) brightness and its [magnetic field strength](#) is known as the bright-magnetic relationship.

The study of the bright-magnetic relationship of sunspots is helpful to understand the brightness and atmospheric structure of sunspots, and provides constraints for the theoretical model of the energy transport mechanism of sunspots.

Previous observational studies on the bright-magnetic relationship of sunspots have mostly focused on stable sunspots, and few have studied the bright-magnetic relationship of decaying sunspots.

The researchers analyzed the difference of the bright-magnetic relationship between the stable and decaying sunspots by using the stray light calibration data from the Solar Dynamics Observatory Helioseismic and Magnetic Imaging.

They found that the difference of the bright-magnetic relationship between stable and decaying sunspots was mainly manifested in the [continuum intensity](#) (brightness) range from 0.35 Iqs to 0.65 Iqs (Iqs is the average continuum intensity of the quiet sun). In the decaying

sunspot, the relationship between the continuum intensity and the transverse magnetic field strength showed a higher degree of dispersion.

"We also found a bifurcation structure in the relationship between the continuum intensity and the transverse magnetic field strength during the formation of the light bridge. This bifurcation structure indicates that the two parts of the umbra separated by the light bridge have different thermodynamic properties," said Prof. Yan.

"The continuum intensity of the sunspot umbra in decaying sunspots is brighter than that of a stable sunspot, and the mean continuum intensity of the sunspot umbra gradually increases during the decay of sunspot. However, the mean continuum intensity of the penumbra does not change with the decay of the sunspot," said Li.

These results indicate that during the decay of sunspot, the temperature of the umbra increases gradually, but the temperature of the penumbra remains constant.

More information: Qiaoling Li et al, Dependence of the Continuum Intensities on the Magnetic Fields at Different Evolution Phases of Sunspots, *The Astrophysical Journal* (2022). [DOI: 10.3847/1538-4357/ac83b3](https://doi.org/10.3847/1538-4357/ac83b3)

Provided by Chinese Academy of Sciences

Citation: Researchers determine bright-magnetic relationship for decaying sunspots (2022, September 26) retrieved 24 May 2024 from <https://phys.org/news/2022-09-bright-magnetic-relationship-sunspots.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.