


Characteristics of older forests can buffer effects of climate change for some bird species

September 6 2022, by Steve Lundeberg



Study area map and photographs of typical vegetation in the study area. (a) Location of the study area, H.J. Andrews Experimental Forest in Oregon. (b) Sampling locations (points; black circles) and altitudinal gradient of the watershed. (c) Vegetation height and stands with harvest history (yellow boundaries). (d) Typical old-growth Douglas-fir-Western hemlock forest and (e)

Douglas-fir second-growth plantation in H.J. Andrews Experimental Forest.
Credit: *Global Change Biology* (2022). DOI: 10.1111/gcb.16353

Old-growth forests and managed forests with old-growth characteristics can provide relief from climate change for some bird species, research by the Oregon State University College of Forestry suggests.

The study led by former Oregon State doctoral student Hankyu Kim builds on earlier research led by co-author Matt Betts, a professor in the Department of Forest Ecosystems and Society, that showed that old forests with big trees and a diversity of tree sizes and species can offer refuge to some types of birds threatened by a warming climate.

The latest findings bear important implications on conservation decisions regarding mature forests, the scientists say, and have even greater relevance because of the new Inflation Reduction Act, which calls for increased resources to map and protect the United States' remaining [old-growth forests](#).

The research, published today in *Global Change Biology*, looked at forest "microclimates." Microclimates are local atmospheric conditions, in areas ranging from a few square meters to many square kilometers, that differ from those of the surrounding area.

Microclimates tend to be most pronounced in areas of rugged and varied topography such as coastal areas, islands and mountains like Oregon's Cascade Range, home to the HJ Andrews Experimental Forest where Kim and Betts did their research.

The OSU scientists and collaborators from Oregon State and the United States Forest Service analyzed eight years of breeding bird abundance

information from an HJ Andrews watershed as well as subcanopy temperature readings and ground- and LiDAR-based vegetation data. They concluded that at locations with cooler microclimates, some [bird species](#) tended to do better—a phenomenon they describe as the "buffering effect."

Some species also fared better in places where the forest had more compositional diversity, referred to as the "insurance effect" because diversity helps ensure the presence of the insects the birds feed on when they most need nutrition and energy—during breeding season.

"To my knowledge, this is the first empirical evidence of any microclimate effect on songbird populations, and of the insurance effect on free-ranging birds," said Kim, now a postdoctoral researcher at the University of Wisconsin-Madison. "Each species may have a slightly different range of thermal optima—the range of thermal conditions they feel comfortable with—and it could be the same for the interaction between forest ecosystems and birds."

Under the current warming regime, he explained, some birds interact with the forest ecosystem to their benefit, while others will find it hard to breed there because the availability of food has changed for the worse.

The scientists found that for five of the 20 bird species they analyzed, abundance trends tended to be either neutral or less negative in cooler microclimates, and the negative effects of warming on two species were reduced in locations with greater forest compositional diversity.

The five species benefiting from the buffering effect were the Swainson's thrush, chestnut-backed chickadee, hermit warbler, varied thrush and Wilson's warbler. The Wilson's warbler and the red crossbill were the two with statistical evidence of benefiting from the insurance

effect.

"If plants leaf out earlier in warm microclimates, causing arthropods to emerge earlier, there is a danger of migratory birds mistiming their breeding with peak food availability," Betts said. "Since leaf-out timing varies by [plant species](#), forests with more plant diversity often have a longer period of insect availability."

The other 14 birds in the analyses were the dark-eyed junco, hermit thrush, McGillivray's warbler, Pacific-slope flycatcher, brown creeper, black-throated gray warbler, golden-crowned kinglet, Hammond's flycatcher, hairy woodpecker, Pacific wren, red-breasted nuthatch, red-breasted sapsucker, western tanager and yellow-rumped warbler.

Seven of the 20 species showed overall declines in abundance over the eight-year study, 2011-18. Nine showed increases and four did display a detectable trend.

"Trends of abundance of five species declined at greater rates in warmer locations than in cooler areas," Kim said. "That suggests microclimates within forested landscapes do provide refugia for those species. Declining species that are sensitive to warm conditions, like the Wilson's warbler, hermit warbler and chestnut-backed chickadee, seemed to benefit the most from refugia effects."

Betts found it interesting that the study led by Kim—whose results Betts said were "collected independently and more rigorously" than those in the research he led in 2017—showed the same species in decline and the same species benefiting from forests with old-growth characteristics.

"The earlier paper was less well done because we didn't measure microclimate directly," Betts said. "Our hypothesis was that [microclimate](#) buffering should work for a high proportion of the

declining species. This current paper is the first time that's been shown."

More information: Hankyu Kim et al, Forest microclimate and composition mediate long-term trends of breeding bird populations, *Global Change Biology* (2022). [DOI: 10.1111/gcb.16353](https://doi.org/10.1111/gcb.16353)

Provided by Oregon State University

Citation: Characteristics of older forests can buffer effects of climate change for some bird species (2022, September 6) retrieved 1 May 2024 from <https://phys.org/news/2022-09-characteristics-older-forests-buffer-effects.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--