


Researchers demonstrate chemically sensitive LiDAR method

September 21 2022



Results of displacement obtained from Fourier transform of the phase modulation of 1D scan of the sample. Credit: Nokia Bell Labs

Researchers have developed a new laser-based technique that can simultaneously perform LiDAR and remote chemical measurements. LiDAR, which stands for light detection and ranging, uses a laser to measure distances, or ranges. Adding chemical information to LiDAR measurements could be useful for applications like remote chemical mapping, detecting trace amounts of chemicals, monitoring industrial processes and quality control.

"By mapping and identifying the composition of the environment, we can augment human interactions and [industrial processes](#) of the future with multi-dimensional object information beyond just ranging and

detection," said Bibek R. Samanta, technical staff at Nokia Bell Labs.

Samanta will present the research at the Frontiers in Optics + Laser Science Conference (FiO LS) meeting being held in Rochester, New York and online 17-20 October 2022. Samanta's presentation is scheduled for Monday, 17 October 2022 October at 12:00 EDT (UTC—04:00).

Combining methods

The new technique, which combines photothermal spectroscopy and LiDAR, resolves [chemical information](#) by detecting sub-nanometer surface deformations due to photothermal absorption of a pump laser. These photothermal effects are caused by intensity modulations of the pump beam.

The researchers used a swept-source laser as a probe beam to perform a LiDAR scan in a frequency-modulated continuous wave configuration. The pump beam was a wavelength-stabilized infrared laser diode laser modulated using a chopper wheel. Both beams were collimated, combined and focused on the same spot about 8 centimeters away. For this setup, the researchers estimated an axial resolution of about 150 microns in air and an imaging depth of about 30 centimeters.

The investigators tested their new approach by using a 3D printed transparent plastic block with 500-micron deep channels containing [epoxy resin](#) mixed with either a green acrylic color or a near-infrared (NIR) absorbing dye. They were able to observe surface displacements of 0.2 to 0.3 nm that resulted from photothermal absorption of the epoxy containing the NIR dye. This agreed with estimated values and was about an order of magnitude greater than the system's baseline noise.

By scanning the sample laterally, the researchers created a chemically

sensitive LiDAR scan that showed the location of the NIR dye but not the green acrylic resin. Although an infrared laser was used in this demonstration, other wavelengths could be used to identify other materials.

"Using tunable [laser](#) systems and a fast-scanning integrated optical assembly, we plan on implementing spectroscopic identification of common household materials to create a 5D map of the environment," Samanta further explains.

Provided by Optica

Citation: Researchers demonstrate chemically sensitive LiDAR method (2022, September 21) retrieved 27 April 2024 from

<https://phys.org/news/2022-09-chemically-sensitive-lidar-method.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--