

As chinook salmon get thinner and fewer, southern resident killer whales struggle to find enough food

September 21 2022, by Fanny Couture



Credit: Unsplash/CC0 Public Domain

As marine species continue to decline worldwide, the southern resident killer whale population—<u>which now stands at 75 individuals</u>—along the



west coast of North America, has baffled scientists who are trying to understand why this population is struggling.

Studied extensively since the mid-1970s

, these whales are considered endangered today.

Food shortage, vessel traffic and water pollutants have affected their fertility, body condition and social behavior. In fact, food shortage is said to be one of the primary causes of the declining whale population, which feed primarily on chinook salmon. This salmon species can constitute up to 90% of the southern resident killer whale's diet during summer.

In the late 1980s, the Canadian government applied <u>restrictive fishing</u> <u>regulations</u> for chinook salmon to protect the population, leading to a <u>drastic decline</u> in <u>commercial fisheries</u>. However, the <u>chinook salmon</u> <u>stocks of the Salish Sea</u> continued to decline. Today, most chinook salmon stocks that the southern resident killer whales feed upon are at very low levels.

As a researcher in ecosystem modeling, I have studied the predator and prey interactions between the southern resident killer whales and their favorite salmon preys since 2019. The declining salmon and whale numbers raised a critical question: Is the southern resident killer whale population solely reliant on the abundance of salmon? And, if so, since when?

The predator and the prey

Along with researchers from the University of British Columbia and marine mammal experts, I developed a <u>bioenergetics model</u> to compare the prey abundance with the energetic requirements of these whales over



the last 40 years.

We estimated that these whales could capture about 12 fish per day and evaluated whether those fish were enough to sustain the energetic requirements of the population from May to October. <u>Chinook salmon populations—targeted by the southern resident killer whales—originate from different geographical areas in the Salish Sea and along the west coast of Vancouver Island.</u>

The energy available to these whales depended not only on the abundance of chinook salmon present in their hunting grounds, but also on the size and age of the fish.

A <u>recent study</u> suggested that most Northeastern Pacific chinook salmon have declined in size over the last decades, and recent <u>evidence points</u> <u>that the older, fatter fish are also becoming more scarce</u> in several populations.

We found that these whales were not getting enough food for six of the last 40 years and lost about 17% of their energetic requirements on average. That is equivalent to an adult human missing out on their daily breakfast.

We found that the southern resident killer whale population also consumed less adult chinook salmon than harbor seals in recent years, and less adult chinook salmon than fisheries over the entire time period. It is now essential to estimate the role of fisheries and marine mammal predation on the chinook salmon populations targeted by these whales.

Who is to blame?

Southern resident killer whales constitute a very small population, making them vulnerable to various factors that could impact their



reproductive ability, social behavior or body growth.

A <u>recent study</u> suggested that the physicality of the southern resident killer whales have been worsening in recent years and that these observations could be linked to a decrease in <u>food availability</u>.

It is crucial to understand the primary causes for the decline in chinook <u>salmon</u> populations as well. Is it a result of predation by other animals, including other marine mammals? Degradation of habitats? Prevalence to diseases? Climate change?

The southern resident killer whales and <u>chinook salmon</u> are two iconic species of the west coast of Canada, both of which are in need of conservation. They have a high economic value for tourism and fisheries and are <u>highly culturally important animals for Coast Salish First Nation</u> <u>Communities</u>.

Predator-prey relationship holds the key

The <u>southern resident killer whales</u> are not like any other killer whales. They have a unique <u>genetic profile</u>, <u>vocalization characteristics</u> and <u>geographical distribution</u>. Preserving such uniqueness in a natural ecosystem is essential for promoting biodiversity.

We need to look at the link between predators and their prey to understand the trends observed in other similar resident killer whale populations. For example, the northern counterpart of the these whales—referred to as the northern resident killer whale population—feed on similar prey species but have been <u>increasing over</u> <u>the last 40 years</u>.

Today, it is unclear how prey drive these two populations in different directions, and future research on northern <u>resident</u> killer <u>whales</u> might



also help us solve the mystery of the decline in the population of its southern counterparts.

It is time for governmental, Indigenous, research and fisheries organizations to unite their efforts to determine what needs to be done in order to protect this whale <u>population</u> and their prey of choice.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: As chinook salmon get thinner and fewer, southern resident killer whales struggle to find enough food (2022, September 21) retrieved 28 June 2024 from <u>https://phys.org/news/2022-09-chinook-salmon-thinner-southern-resident.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.