


Chiral quasi bound states in the continuum for a high-purity circularly polarized light source

September 8 2022



High purity circularly polarized spontaneous emission and lasing from the resonant metasurface with near-unity intrinsic chirality. Credit: Xudong Zhang

An ultracompact circularly polarized light source is a crucial component for the applications of classical and quantum optics information processing. The development of this field relies on the advances of two fields: quantum materials and chiral optical cavities. Conventional approaches for circularly polarized photoluminescence suffer from incoherent broadband emission, limited DOP, and large radiating angles. Their practical applications are constrained by low efficiency and energy waste to undesired handedness and emission directions. The chiral microlasers can have large DOPs and directional output, but only in specific power ranges. Most importantly, their subthreshold performances plummet significantly. Up to now, the strategy for simultaneous control of chiral spontaneous emission and chiral lasing is still absent.

In a new paper published in *Science*, researchers from Harbin Institute of Technology and Australian National University employ the physics of chiral quasi [bound states](#) in the continuum (BICs) and demonstrate the efficient and controllable emission of circularly polarized light from resonant metasurfaces.

BICs with integer topological charge in momentum space and a theoretically infinite Q factor have been explored for many applications including nonlinear optics and lasing. By introducing in-plane asymmetry, BICs turn to be quasi-BICs with finite but still high Q factors. Interestingly, the integer topological charge of BICs mode would split into two half integer charges, which symmetrically distribute in momentum space and correspond to left- and right-handed circular polarization states, also known as C points.

At the C points, [incident light](#) with one circular polarization state can be coupled into the nanostructures and produce dramatically enhanced local

electromagnetic fields. The other polarization state is decoupled and almost perfectly transmit. Such characteristics are well known but rarely applied to light emissions. "This is mainly because the C points usually deviate from the bottom of band. They have relatively low Q factor and cannot be excited for lasing actions," says Zhang.

To realize the chiral light emission, a key step is to combine the local density of states with the intrinsic chirality at C points. If one C point is shifted to the bottom of the band, the Q factor of the corresponding chiral quasi-BIC can be maximal. According to the Fermi's golden rule, the radiation rate of one circularly polarized spontaneous emission is enhanced, whereas the other polarization is inhibited. Both the Q factor and the radiation rate reduces dramatically with the emission angle. As a result, high-purity and highly directional light emission can be expected near the Γ point.

"Of course, the other C point can support similar high chirality with opposite handedness. However, that point also deviates from the maximal Q factor and less be enhanced. Therefore, our metasurface only produces one near unity circular polarization with high directionality around the normal direction," says Zhang.

The control of C points in momentum space closely relates to the maximization of chirality in normal direction. In principle, the realization of chirality relates to the simultaneous breaking of in-plane and out-of-plane mirror reflection symmetries. In this research, the team has introduced an out-of-plane asymmetry, the tilt of nanostructures. For an in-plane asymmetry, there is one out-of-plane asymmetry that can move one C point to Γ point. "We find two types of asymmetries are linearly dependent on one another. This makes the optimization of chirality in normal direction very easy," says Zhang.

In experiments, the researchers have fabricated the metasurfaces with a

one-step slanted reactive ion etching process and characterized the emissions. Under the excitation of a nanosecond laser, they have successfully demonstrated the chiral emissions with a DOP of 0.98 and a far field divergent angle of 1.06 degree. "Our circularly [light source](#) is realized with the control of C point in momentum space and local density of state. It is independent of the excitation power," says Zhang. "This is the reason that we can achieve the high Q, high directionality, and high purity [circular polarization](#) emission from spontaneous emission to lasing."

Compared to conventional approaches, the chiral quasi-BIC provides a way to simultaneously modify and control spectra, radiation patterns, and spin angular momentum of photoluminescence and lasing without any spin injection. This approach may improve the design of current sources of chiral light and boost their applications in photonic and quantum systems.

More information: Xudong Zhang et al, Chiral emission from resonant metasurfaces, *Science* (2022). DOI: [10.1126/science.abq7870](https://doi.org/10.1126/science.abq7870). www.science.org/doi/10.1126/science.abq7870

Provided by Harbin Institute of Technology

Citation: Chiral quasi bound states in the continuum for a high-purity circularly polarized light source (2022, September 8) retrieved 26 June 2024 from <https://phys.org/news/2022-09-chiral-quasi-bound-states-continuum.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.