

Crime-scene technique identifies asteroid sites

September 1 2022



Location, age, and diameter of confirmed impact craters: (A) Kaali Main and Kaali 2/8, both in Estonia; (B) Morasko in Poland; and (C) Whitecourt in Canada, along with field images of in situ charcoals (at thin white arrows) found within their proximal ejecta blankets. Charcoals are present throughout the ejecta (panels A1, B1, and B2) but are most abundant close to the base (panels



A2, C1, and C2). Thick white arrows in panels A and B point out the locations of trenches where charcoals visible in A1, A2, B1, and B2 were collected. Numbers in C and in the field photos refer to sample numbers. Credit: *Geology* (2022). DOI: 10.1130/G50056.1

Tens of tons of extraterrestrial solid material collide with Earth daily. Most of this material is small enough that it burns up in the atmosphere, but some fragments are large enough to cause quite a predicament. In 2013, a 20-meter-diameter body exploded over Chelyabinsk, seriously injuring more than 1,500 people.

The most recent impact crater on Earth was formed in 2007, when an asteroid crashed into a small village in Peru. A 1947 asteroid impact in far-eastern Russia resulted in the formation of the youngest crater-strewn field on our planet: Sikhote-Alin. The most impressive, relatively recent, extra-terrestrial event happened in 1908, when a body exploded over Siberia, flattening 2000 km² of forest.

We can prepare for this natural hazard only if we understand how often these kinds of small impacts happened in the past and how they influenced the environment.

A new article in the Geological Society of America's journal *Geology* shows that analyzing bodies of organisms killed by an impact of asteroids can teach us exactly how much damage occurs at the spot of such a cosmic collision. The research team dug out trenches in rims of four craters (Kaali Main and Kaali 2/8 in Estonia, Morasko, in Poland, and Whitecourt in Alberta, Canada), located on two different continents that formed thousands of years apart.

Dr. Jüri Plado and Dr. Argo Jõeleht noted that "surprisingly, in all those



places we found the same thing: millimeter- to centimeter-sized pieces of charcoal intermixed within material ejected during its formation and located at the same place in respect to the crater."

Dr. Ania Losiak, the lead author of this study, from the Institute of Geological Sciences, Polish Academy of Sciences, and the University of Exeter, said, "At first we thought those charcoals were formed by wildfires that occurred shortly before the impact, and charcoals just got tangled in this extraterrestrial situation.

"But something was not right with this hypothesis—there were too many coincidences; why would there be large wildfires shortly before formation of four different small impact craters divided by thousands of kilometers and years? Why would it be found only in a very specific location within the proximal ejecta blanket? It made no sense, so we decided to investigate further and analyze properties of charcoal pieces found intermixed within material ejected from craters and compare it with wildfire charcoals."

Like bodies studied in a criminal investigation, the properties of organic remnants turned into charcoal reflect the conditions in which they were killed. Based on their properties, we can clearly recognize charcoals formed as a result of a wildfire and those found within proximal ejecta of impact craters.

Professor Claire Belcher from the University of Exeter explained, "Impact charcoals are really weird: They were all formed in much lower temperatures than wildfire charcoals, they lack sections that were formed while directly touching the flame, and they are all very similar to each other, while in a fire it is common to find strongly charred wood just next to barely affected branches."

"This is definitely not what we expected when we started this study: We



think that impact charcoals were formed when fragments of trees shattered by the impact were intermixed with local material ejected from the crater," added Losiak.



Digital elevation model of Kaali craters. Credit: Argo Jõeleht





Looking for charcoal. Credit: Juri Plado





Finding charcoal within Kaali 2-8 crater ejecta. Credit: Juri Plado





Charcoal with wood structure preserved on which it is possible to recognize the type of a tree killed by an asteroid. Image under Scanning Electron Microscope. Credit: Ania Losiak





Fragment of organism killed and baked alive by an asteroid (picture under optical microscope). Credit: Ania Losiak

"This study improves our understanding of environmental effects of small impact crater formation so that in the future, when we discover an asteroid a few meters across or more coming our way only a couple of weeks before the impact, we will be able to more precisely determine the size and type of evacuation zone necessary," said Professor Chris Herd from the University of Alberta.

"Our research may also help to find new impact craters on Earth; we expect that we are missing from our records more than ten craters



formed within the last ten thousand years. We need to find them, before their relatives visit us unexpectedly," explained professor Witek Szczuciński from Adam Mickiewicz University in Poznan.

More information: A. Losiak et al, Small impact cratering processes produce distinctive charcoal assemblages, *Geology* (2022). <u>DOI:</u> <u>10.1130/G50056.1</u>

Provided by Geological Society of America

Citation: Crime-scene technique identifies asteroid sites (2022, September 1) retrieved 29 June 2024 from <u>https://phys.org/news/2022-09-crime-scene-technique-asteroid-sites.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.