


Differential impacts of adult trees on offspring and non-offspring recruits in a subtropical forest

September 16 2022



The figures are the observed and expected sapling recruitment distribution estimated from the seedling distribution (A) and juvenile recruitment distribution estimated from the sapling distribution (B). These show that the peak recruitment distances of offspring shift away from mother trees over life stages, implying a higher mortality rate for seedlings recruiting near their mother trees. Credit: Science China Press


An important mechanism promoting species coexistence is conspecific negative density dependence (CNDD), which inhibits conspecific neighbors by accumulating host-specific enemies near adult trees and thus promoting species coexistence by freeing up space for heterospecific species.

Natural enemies may be genotype-specific. Whether within-species genetic relatedness between seedlings and adult neighbors regulates the strength of CNDD is one of the keys explaining the mechanism of species coexistence. However, this remains largely unexplored in natural forests due to the difficulty in assessing the parent-offspring relationship between individuals within [natural populations](#).

Recently, a study from Dr. Keping Ma group, Institute of Botany, Chinese Academy of Sciences was conducted in a 24-ha subtropical evergreen broad-leaved forest dynamics plot. 3,002 individuals of the most dominant tree species *Castanopsis eyrei* were sampled and 12 [microsatellite markers](#) were used to identify the parent-offspring relationship to estimate the effects of adult-seedling genetic relatedness on seedling recruitment.

The results show that 1) peak recruitment distances of offspring shift

away from mother trees over life stages. Offspring have significantly lower recruitment efficiencies in the vicinity of mother trees. 2) Recruitment efficiency (proxy of survival probability) of offspring compared with non-offspring near adult trees during the seedling-sapling transition, suggesting genotype-dependent interactions drive tree demographic dynamics. 3) The genetic similarity between individuals of same cohort decreased in late life history stages, indicating genetic-relatedness-dependent tree mortality throughout ontogeny.



The figures show the recruitment efficiencies (RE, the recruit ratio of early life stage to the later) of offspring and nonoffspring during seedling–sapling (A) and sapling–juvenile (B) transitions along the distance from the focal trees for *Castanopsis eyrei*. These suggest higher survival chances for offspring when recruit far from their mother trees during seedling–sapling transition. Credit: Science China Press

Overall, the results indicate that the strength of CNDD and population dynamics depend on both [genetic relatedness](#) and spatial distance to conspecific adults, implying genotype-specific [natural enemies](#) may be a key driver. Further research on the interaction between genotype-specific enemies and their [host plants](#) is essential to fully understand the underlying mechanisms of CNDD.

The research was published in *Science China Life Sciences*.

More information: Fang Wang et al, Differential impacts of adult trees on offspring and non-offspring recruits in a subtropical forest, *Science China Life Sciences* (2022). [DOI: 10.1007/s11427-021-2148-7](https://doi.org/10.1007/s11427-021-2148-7)

Provided by Science China Press

Citation: Differential impacts of adult trees on offspring and non-offspring recruits in a subtropical forest (2022, September 16) retrieved 4 May 2024 from <https://phys.org/news/2022-09-differential-impacts-adult-trees-offspring.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--