

## **Dual-lattice Kerker effects: controlling light scattering with incident polarization and angle**

September 8 2022, by Li Yuan



(a)(b) Schematics of periodic silicon nanodisks under oblique illumination of sor p-polarized plane wave. ED-SLRorMD-SLR arises from diffractive coupling (indicated by the luminous stripes) of localized Mie EDR or MDR in individual nanodisks, asindicated by blue or red double-headed arrows. The spectral overlap of ED-SLR and MD-SLR causes resonant lattice Kerkereffect.(c)–(f) Simulated angular-resolved zeroth-order (c)(d) reflectance and (e)(f) transmittance spectra for (c)(e) s-polarizationand(d)(f) p-polarization. Red circles indicate the occurrence of the resonant lattice Kerker effect. Credit: *Nano Research*. DOI: 10.1007/s12274-022-4988-9



By using all-dielectric nanostructures, light can be scattered in a welldefined direction, which is the so-called generalized Kerker effects. These effects, however, are usually polarization-independent or realized for a particular polarization only.

A research group led by Dr. Li Guangyuan from the Shenzhen Institute of Advanced Technology (SIAT), Chinese Academy of Sciences (CAS), proposed and experimentally demonstrated polarization-controlled dual-<u>lattice</u> Kerker effects in periodic silicon nanorods.

These results enable active tuning of the Kerker effects by varying the incident polarization or angle, and can be used in various applications including the manipulation of the direction, polarization, and phase of the scattered light, which are essential in nanophotonic chips.

This study was published in Nano Research.

In the dual-lattice Kerker effects, the incident angles (zero reflection and unitary transmission), which are referred to as the lattice Kerker angles, can be the same or different for the s- and p-polarizations, depending on the choice of the silicon nanodisks' diameter and height. These lattice Kerker angles can be further tuned within large ranges by varying the lattice periods in both directions.

Other generalized Kerker effects reported in the literature mainly work under normal incidence, and are realized by varying the geometry parameters. This requires stringent choice of parameters and very careful fabrication. "By introducing the lattice effect, the so-called lattice Kerker effect can be realized through varying the incident <u>angle</u>. This merit enables active tuning of the Kerker effect in an as-fabricated sample, and thus significantly facilitates the design and fabrication," said Dr. Li.



An unexpected phenomenon is that, high-order multipoles such as electric and magnetic quadrupoles are also involved and become important when the periodic silicon nanodisks are obliquely illuminated by p-polarized light. This results in different dispersion relationships between the electric dipole surface lattice resonance (ED-SLR) under ppolarization and the magnetic dipole surface lattice resonance (MD-SLR) under s-polarization, leading to different lattice Kerker angles for the s- and p-polarizations.

**More information:** L. Xiong et al, Polarization-controlled dual resonant lattice Kerker effects. *Nano Research*. DOI: <u>10.1007/s12274-022-4988-9</u>. <u>www.sciopen.com/article/pdf/10 ...</u> <u>12274-022-4988-9.pdf</u>

Provided by Chinese Academy of Sciences

Citation: Dual-lattice Kerker effects: controlling light scattering with incident polarization and angle (2022, September 8) retrieved 28 June 2024 from <u>https://phys.org/news/2022-09-dual-lattice-kerker-effects-incident-polarization.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.