

Efficient species monitoring method proves its worth for conservation management

September 20 2022



An image representative of pāua beds off the coast of Warrington, Otago, New Zealand. (a) Blackfoot pāua (*Haliotis iris*) beds spread across rocks in a patchily distributed population. (b) three individual Blackfoot pāua covered in sand.

Credit: Dr. Matthew Desmond (matthew.desmond@otago.ac.nz); *Environmental DNA* (2022). DOI: 10.1002/edn3.352

Genetic information is a critical part of conservation research but has proven invasive and costly—until now.

A University of Otago study compared traditional tissue sampling with environmental DNA sampling to obtain population genetic information. It showed there is potential for this new approach to help with [conservation management](#).

eDNA is an effective tool to monitor the presence of species in water—as an animal swims by in the ocean, it releases DNA into the environment through excrement, mucus, or the shedding of scales which can be collected and analyzed by researchers.

Study lead author Dr. Clare Adams, of the Coastal People: Southern Skies Center of Research Excellence, says the group wanted to push this method one step further, obtaining genetic information on a population, specifically blackfoot pāua.

"We wanted to determine how well eDNA methodologies could be used to monitor populations of marine species.

"Conservation research has shown that [genetic information](#) is useful to support management decisions, but it is difficult to come by and involves invasive sampling of the very species you are hoping to protect and manage.

"eDNA methods are increasingly cheaper, easier, and less time consuming than traditional genetic sampling, so they could be used to help monitor how genetically healthy and biodiverse populations of target organisms may be," she says.

The study, published in *Environmental DNA*, took tissue samples of pāua and [water samples](#) from near the pāua colonies.

Dr. Adams says eDNA proved to be an effective method in analyzing common genetic variations, though rare genetic variants (present in less than 5% of the population) were more difficult to detect.

The researchers focused on pāua due to its cultural importance to Māori and commercial importance to the global fisheries industry.

"Genetic diversity is essential to monitor because it can tell us about how inbred pāua stocks might be, as well as hinting at [gene flow](#), and ultimately can highlight mechanisms of evolution.

"We hope some of the work done here can help communities monitor their pāua stock."

More information: Clare I. M. Adams et al, Environmental DNA reflects common haplotypic variation, *Environmental DNA* (2022). [DOI: 10.1002/edn3.352](https://doi.org/10.1002/edn3.352)

Provided by University of Otago

Citation: Efficient species monitoring method proves its worth for conservation management (2022, September 20) retrieved 18 June 2024 from <https://phys.org/news/2022-09-efficient-species-method-worth.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.