

Lattice distortion of perovskite quantum dots induces coherent quantum beating

September 9 2022



Lattice distortion in lead halide perovskite quantum dots leads to a fine structure gap and coherent exciton quantum beating. Credit: DICP

A research group led by Prof. Wu Kaifeng from the Dalian Institute of Chemical Physics (DICP) of the Chinese Academy of Sciences (CAS), in collaboration with Dr. Peter C. Sercel from the Center for Hybrid Organic Inorganic Semiconductors for Energy, recently reported the utilization of lattice distortion in lead halide perovskite quantum dots (QDs) to control their exciton fine structure.

The study was published in *Nature Materials* on Sept. 8.



It is well known that shape or crystal anisotropy in QDs, which are tiny semiconductor nanoparticles, results in energy splitting of their optically bright excitons (bound <u>electron-hole pairs</u>), known as fine structure splitting (FSS). These excitons form an important playground for quantum information science. For example, the excitons' FSS can be exploited for coherent control of quantum states for <u>quantum computing</u>, or for polarization-entangled photon-pairs in <u>quantum optics</u>, although for the latter it is important to suppress the magnitude of splitting.

Traditionally, studying FSS usually requires single or just a few QDs at liquid-helium temperature, because of its sensitivity to QD size and shape. Measuring FSS at an ensemble-level, let alone controlling it, seems impossible unless all the dots are made to be nearly identical.

In this study, by using ensemble-level femtosecond polarized transient absorption, the researchers observed clear bright-exciton FSS in solution-processed CsPbI₃ perovskite QDs, which is manifested as exciton quantum beats (periodic oscillations of kinetic traces).

"Even more amazingly, the beat frequency, as determined by the FSS energy, of a given sample can be continuously controlled by changing the temperature. This is an unprecedented result, meaning that now scientists can facilely control FSS through temperature," said Prof. Wu.

The researchers also found that the temperature-dependent FSS was related to the interesting, highly-dynamic lattice of lead halide perovskites. Lowering the temperature led to a more distorted leadiodide octahedral framework.

Calculations indicated that, because these orthorhombic-phase QDs were actually still bounded by the pseudocubic family of crystal planes, the lattice distortion results in an avoided crossing fine-structure gap between bright exciton. This gap was responsible for the observed FSS,



and it could be detected in spite of QD size and shape heterogeneity across an ensemble sample.

"Lattice distortion in CsPbI₃ perovskites is well known in the photovoltaic community, as it is connected to the issue of phase stability of perovskite solar cells, but nobody has previously connected it experimentally to the exciton fine structure" said Prof. Wu. "Our study demonstrates that this material property can actually be harnessed to control the bright-<u>exciton</u> splitting in quantum dots for quantum information technologies."

More information: Yaoyao Han et al, Lattice distortion inducing exciton splitting and coherent quantum beating in CsPbI3 perovskite quantum dots, *Nature Materials* (2022). DOI: 10.1038/s41563-022-01349-4

Provided by Chinese Academy of Sciences

Citation: Lattice distortion of perovskite quantum dots induces coherent quantum beating (2022, September 9) retrieved 2 May 2024 from <u>https://phys.org/news/2022-09-lattice-distortion-perovskite-quantum-dots.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.