

## Microplastics are common in homes across 29 countries: New research shows who's most at risk

September 6 2022, by Mark Patrick Taylor, Neda Sharifi Soltani and Scott P. Wilson



Credit: Pexels

The evidence is clear: microplastics have contaminated <u>every corner of</u> <u>the globe</u>. We can't escape exposure to these tiny bits of plastic (less than 5mm across) in the environment, which includes the homes where people <u>spend most of their time</u>.



Recent research has discovered <u>microplastics in the blood of humans</u>. However, the question of harm to humans <u>remains unresolved</u>. Despite concerns that some substances in microplastics could cause cancer or damage our DNA, we still have a poor grasp of the true risks of harm.

Our <u>study of global microplastics</u> exposure inside homes across 29 countries, published today, shows people living in lower-income countries and young children are at greater risk of exposure. But our analysis of the chemical composition of microplastics in the home shows the specific health risk is surprisingly low. The study covered all the continents, including Australia.

The current challenge in understanding health risks from microplastics is the very limited data on toxic effects of the petrochemicals used in plastics production.

A recurrent theme in the environmental health research literature is that early concerns about suspect chemicals and related compounds, including those found in plastics, <u>were eventually justified</u>. The effects of suspect substances only become clear after extensive toxicological and epidemiological research.

Microplastics are "ubiquitous" and have been detected in fresh water, wastewaster, food, air and drinking water.

But a new WHO study says more research is needed to firmly conclude that microplastics pose a risk to human health.<u>https://t.co/3T6ngvs6q2</u>

— NPR (@NPR) <u>August 23, 2019</u>

### What did the new study look at?



Our study investigated three main questions relating to exposure to microplastics inside homes:

- What are the impacts in different countries across the world?
- Who is most at risk?
- What are the specific <u>health risks</u>?

We reached out to residents across 29 countries to collect their indoor atmospheric dust over a one-month period. At 108 homes sampled across these countries, we also collected information about households and behaviors. This helped us to better understand possible sources and causes of microplastics in dust. These data included:

- how often floors were cleaned
- flooring type
- presence or absence of children
- number of people living in each home
- percentage of full-time workers.



# Sources of microplastics in homes



Author provided, The Conversation

In each home, atmospheric dust particles were collected in specially cleaned and prepared glass Petri dishes. We measured the levels of microplastics in the collected dust using a suite of microscopic techniques and instruments. We used <u>infrared spectroscopy</u>—which identifies substances by how they interact with light—to determine the chemical composition of these microplastics.

#### What did the study find?

The household dust contained a wide variety of synthetic <u>polymer</u> fibers. The most common were:



- polyester (as <u>polyethylene terephthalate</u>) at 9.1%, which is used in clothing fabrics
- polyamide (7.7%), which is mainly used in textiles
- polyvinyls (5.8%), which are used in floor varnishes
- polyurethane (4.4%), which is used in surface coatings of furniture and in bedding
- polyethylene (3.6%), a common polymer used in food containers and reusable bags.

We examined the prevalence and risk of microplastics according to the gross national income of each country, grouped as low, medium and high-income (such as Australia). Overall, we found lower-income countries have higher loads of microplastics, which were deposited at an average daily rate of 3,518 fibers per square meter. The rates for medium-income and high-income countries were 1,268 and 1,257 fibers/m<sup>2</sup>/day.

In low-income countries, the most prevalent synthetic polymer fibers were made of polyurethane (11.1% of all fibers in samples). In high-income countries, polyamide and polyester were the most prevalent <u>microplastic</u> types (11.2% and 9.8% respectively).

### So what are the health risks?

For the first time we could attribute the health risk across countries according to incomes. Our analyses showed lower-income countries are at higher risk from microplastic pollution. This aligns with <u>research</u> findings on other toxic exposures—poorer countries and people are most at risk from pollution.

Nevertheless, we found the overall risk from microplastics exposure was low. We used the US Environmental Protection Agency's <u>toxicity</u> <u>information</u> on polymers in the microplastics to calculate health risk based on the types and levels we detected.



#### Surveyed countries and most common fibre types

Three categories of countries based on 2021 Gross National Income, according to the World Bank, with the most prevalent fibre types and synthetic polymers for each category.



	Most prevalent fibre types	Most prevalent synthetic polymer
High-income countries	Synthetic polymers 46%	Polyamide
Medium-income countries	Natural fibres 43%	Polyester
Low-income countries	Synthtic polymers 39%	Polyurethane

Credit: The Conversation

Low-income countries had a higher lifetime risk of cancers due to indoor microplastic exposure at 4.7 people per million. High-income countries were next at 1.9 per million, with medium-income countries at 1.2 per million.



We attributed these differences in cancer risk to the different percentages of carcinogenic substances in the microplastics found in household dust.

We calculated the sum of the carcinogenic risk from inhalation and ingestion of the following chemicals in the microplastic fibers: vinyl chloride (polyvinyl chloride), acrylonitrile (polyacrylics) and propylene oxide (polyurethane). Because toxicity data for polymers are limited, the assessment was a minimum estimate of true risk.

Children are at greater risk irrespective of income, which is true for many types of environmental exposures. This is because of their smaller size and weight, and tendency to have more contact with the floor and to put their hands in their mouths more often than adults.

Our analysis indicated that the microplastics came mainly from sources inside the home, and not from outside. Synthetic polymer-based materials are used widely in high-income countries in products such as carpets, furniture, clothing and food containers. We anticipated levels of microplastic shedding in the home might be greater in these countries.

However, analysis of the data showed the only factor obviously linked with levels of microplastics in deposited dust was how often they were vacuumed. Frequent vacuuming reduces microplastic levels.

Vacuuming was more frequent in higher-income <u>countries</u>. Factors that influence the type of cleaning include people's preference for sweeping and mopping versus vacuuming, as well as their access to and capacity to afford electronic vacuum cleaners.

### What can we do to reduce the risks?

Based on this and our previous study data, it is clear vacuuming



regularly, instead of sweeping, is associated with less airborne microplastics indoors. Other obvious actions—such as choosing natural fibers for clothing, carpets and furnishings instead of petrochemical-based polymer fibers—can reduce the shedding of microplastics indoors.

Future research needs to focus on developing more complete profiles of the harmful effects of each of the toxic petrochemical-based synthetic polymers that can produce microplastics. This will give us a better understanding of the risks of exposure to these ubiquitous pollutants.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Microplastics are common in homes across 29 countries: New research shows who's most at risk (2022, September 6) retrieved 11 August 2024 from <a href="https://phys.org/news/2022-09-microplastics-common-homes-countries.html">https://phys.org/news/2022-09-microplastics-common-homes-countries.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.