


# The process of waves carrying plasma heat is observed for the first time

September 28 2022



Heat is carried by electromagnetic waves, which simultaneously slow down the high-speed particle beam (red line) and heat up the plasma particles (green line). Distortion of the velocity distribution is observed due to the acceleration of the particles as a result of the electromagnetic waves. Credit: National Institute for Fusion Science

In fusion power generation, it is essential that the high-energy particles generated by a fusion reaction in hot plasma heat it to sustain further fusion reactions. The key to this self-heating of the plasma is whether it can be heated by waves created by the high-energy particles.

A research group led by Professor Katsumi Ida, Assistant Professors Tatsuya Kobayashi and Mikiro Yoshinuma of the National Institute for Fusion Science and Professor Yuto Kato of Tohoku University, has measured the time variation of the plasma velocity profile in the Large Helical Device (LHD) at the National Institute for Fusion Science and found that [electromagnetic waves](#) produced by high-energy particles carry heat through a process called Landau damping. This is the first observation in the world of this process. A paper summarizing the results of this research was published in *Communications Physics* on Sep. 28.

Until now there has been no method to directly measure the plasma heating process caused by electromagnetic waves generated inside the plasma, so it has not been known whether this process actually exists. In order to capture it, Professor Katsumi Ida and his research group have worked to develop a new measurement system.

In order to directly measure the heating process, it is necessary to determine the time variation of the velocity distribution, which indicates which velocity particles are present and in what proportion. To this end, they injected high-speed atoms into the plasma and used a method to measure the velocity distribution of plasma particles at high speed from the wavelength distribution of light emitted from the plasma (high-speed charge exchange spectroscopy). Professor Ida and his colleagues took on the challenge of ultrahigh-speed measurement, which had been considered difficult, and succeeded in measuring the time variation of the velocity distribution of plasma particles at an ultrahigh-speed of 10

kHz (10,000 times per second).

In the LHD, experiments are being conducted to investigate plasma self-heating, using a high-speed particle beam, that simulates high-energy particles from nuclear fusion reactions. In this experiment to simulate self-heating, a newly developed measurement system was used to measure in detail the time variation of the velocity distribution of plasma particles. As a result, it was discovered for the first time in the world that the plasma is heated due to the slowing down of the high-speed particle beam and the distortion of the velocity profile of the plasma particles caused by the generation of electromagnetic waves inside the plasma.

The reason for this distortion of the [velocity](#) profile was found to be that the energy of the high-speed particle beam was transferred to the electromagnetic wave through a process called Landau damping, and the energy of the electromagnetic wave was transferred to the plasma particles. In other words they observed that the electromagnetic waves carried the energy of the high-speed particle beam to the plasma and heated it.

Professor Ida said, "For self-heating of plasma in fusion power generation, it is not enough for [high-energy particles](#) to collide with plasma ones and heat them, so heating by other processes is also necessary. This result, which shows that electromagnetic waves generated inside the [plasma](#) can heat it, provides important knowledge for [fusion](#) research. Furthermore, it will contribute to the study of the Earth's magnetosphere, where particle acceleration occurs by a similar process, and will promote future interdisciplinary research."

**More information:** Katsumi Ida et al, Direct observation of mass-dependent collisionless energy transfer via Landau and transit-time damping, *Communications Physics* (2022). [DOI: 10.1038/s42005-022-01008-9](#)

Provided by National Institutes of Natural Sciences

Citation: The process of waves carrying plasma heat is observed for the first time (2022, September 28) retrieved 2 May 2024 from <https://phys.org/news/2022-09-plasma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.