

## Studying predatory behavior in the bacterial kingdom

September 14 2022, by Meike Drießen





Graphical abstract. Credit: *Cell Reports* (2022). DOI: 10.1016/j.celrep.2022.111340

Bacteria have a variety of survival strategies to ensure a sufficient food supply in their densely populated habitats. Certain species of bacteria kill microorganisms of another species, decompose their cells and absorb them as nutrients. The exact mechanism of this process is largely unknown.

A research team from the Chair of Microbial Biology headed by Dr. Christine Kaimer has examined these processes in depth. Together with colleagues from the U.S., the researchers from Ruhr-Universität Bochum (RUB) present their findings in the journal *Cell Reports*.

## **Stop on contact**

Little has been known to date about the relationship between predators and prey in the bacterial kingdom. But researchers suspect that bacterial predators can effect significant changes in the composition of a microbiome and thus shape the ecology of their habitat. To learn more about bacterial predator-prey relationships, Christine Kaimer's team studied the predatory bacterium Myxococcus xanthus, which is commonly found in soil. Recent work has revealed that M. xanthus kills its prey cell in direct cell-cell contact: the predator approaches a prey cell, stops once contact has been made, and then induces cell death and decomposition within a few minutes. The researchers have analyzed the molecular mechanisms of this process in close detail.

"We made specific genetic modifications to M. xanthus and then monitored the interaction of predator and prey <u>cells</u> under the microscope in real time," explains Christine Kaimer. "As a result, we



were able to show that two highly specialized <u>protein</u> secretion systems work together in the killing process. These are protein complexes that presumably transport proteins from the predator cell into the prey cell." Initially, a so-called Tad-like apparatus induces <u>cell death</u> to prey. A type 3-like system is then needed for decomposing the prey cell so that the nutrients can subsequently be absorbed.

## Killing initiated with pinpoint accuracy

Images with fluorescently labeled proteins showed that the protein secretion systems accumulate in the M. xanthus cell exactly at the contact site between predator and prey. It appears that M. xanthus senses contact with a prey cell and can initiate the killing process with great precision.

"We don't yet understand how exactly the protein secretion systems react to prey contact and which potentially toxic proteins they release," says Christine Kaimer. "In any case, bacterial predation behavior is a fascinating field of research, where we still have a lot to learn about how bacteria coexist in their natural environment."

**More information:** Susanne Thiery et al, The predatory soil bacterium Myxococcus xanthus combines a Tad- and an atypical type 3-like protein secretion system to kill bacterial cells, *Cell Reports* (2022). DOI: 10.1016/j.celrep.2022.111340

## Provided by Ruhr-Universitaet-Bochum

Citation: Studying predatory behavior in the bacterial kingdom (2022, September 14) retrieved 3 May 2024 from <u>https://phys.org/news/2022-09-predatory-behavior-bacterial-kingdom.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.