

# For the first time, robots on Mars found meteorite impact craters by sensing seismic shock waves

September 20 2022, by Katarina Miljkovic



Credit: NASA / JPL-Caltech

Since 2018, NASA's InSight mission to Mars has recorded seismic waves from more than 1,300 marsquakes in its quest to probe the internal structure of the red planet. The solar panels of the car-sized robotic lander have become caked with Martian dust, and NASA scientists expect it will completely power down by the end of 2022.



But the internal rumblings of our planetary neighbor aren't the only things that InSight's seismometers detect: they also pick up the thuds of <u>space</u> rocks crashing into the Martian soil.

In <u>new research</u> published in *Nature Geoscience*, we used data from InSight to detect and locate four high-speed meteoroid collisions, and then tracked down the resulting craters in <u>satellite images</u> from NASA's Mars Reconnaissance Orbiter.

#### **Rocks from space**

The solar system is full of relatively small rocks called meteoroids, and it's common for them to collide with planets. When a meteoroid encounters a planet with an atmosphere, it heats up due to friction—and may burn up entirely before reaching the ground.

On Earth, we know these incoming meteoroids as shooting stars, or meteors: beautiful events to observe in the night sky. Sometimes a meteoroid explodes when it reaches the thicker atmosphere closer to the ground, creating a spectacular airburst.

Occasionally, a space rock survives its fiery path through the air and drops to the ground, where it is known as a meteorite.

A few of these meteorites hit the surface at such speed they blast a hole in the ground called an impact <u>crater</u>. Compared to a human lifetime, these events are very rare on Earth.

## **Recording space rock impacts**

Scientists have detected the vibrations from meteoroid airbursts using seismic detectors numerous times, including <u>a recent survey</u> of bright



meteors above Australia.

However, only once has a high-speed space rock crashing into the ground been observed both visually and with modern seismic equipment. This was an <u>impact crater</u> that <u>formed in 2007</u> near the village of Carancas in Peru.

Numerous impacts were detected on the moon by the network of seismic sensors set up during the US Apollo missions of the 1960s and '70s. However, there was no recording of a natural impact associated with visual detection of a new crater.

The closest things to such an observation were artificial impacts: the crash-landings of the booster rockets of the ascent modules that lifted Apollo astronauts off the moon.

These human-made impacts on the moon were recorded both in <u>seismic</u> <u>data</u> and visual imagery from orbit. These data were <u>recently used</u> to test simulations of how impacts produce seismic waves.

## **Martian meteorites**

Incoming meteoroids make waves in the atmosphere and also the ground. The atmosphere of Mars is equivalent to 1% of the Earth's, and has a different chemical composition. This means meteor events on Mars take a different form.

For meteor events large enough to drop a meteorite, the fate of the meteorite and any resulting crater is <u>different</u> from what we have come to expect on our home planet.

Here on Earth, or on the moon, single craters are the norm. On Mars, however, about half the time a high-speed space rock will burst in the



atmosphere shortly before impact, resulting in a tightly grouped cluster of craters.

The separation of these individual fragments remains close at ground level, forming a <u>cluster of small impacts</u>.

#### From vibrations to craters

Recently, the InSight mission has observed acoustic and seismic waves from four <u>meteoroid</u> impact events. These waves travel at different speeds, and comparing their different arrival times and other properties allowed us to estimate the location of the impacts.

These impact locations were then confirmed with satellite imaging from the Mars Reconnaissance Orbiter.

Knowing the size and exact location of these impact craters helps us calculate the size and speed of the incoming space <u>rock</u> and how much energy the impact released.

Once we are confident we know something about the impact that created the seismic waves we detected, we can use the waves to learn about the interior of Mars. What's more, when we compare seismic observations on Mars with observations from Earth and the moon, we can learn more about how the planets formed and how the solar system evolved.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: For the first time, robots on Mars found meteorite impact craters by sensing seismic



shock waves (2022, September 20) retrieved 24 May 2024 from https://phys.org/news/2022-09-robots-mars-meteorite-impact-craters.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.