

New seafloor images off the West Antarctic Ice Sheet upend understanding of Thwaites Glacier retreat

September 5 2022



The R/V Nathaniel B. Palmer photographed from a drone at Thwaites Glacier ice front in February 2019. Credit: Alexandra Mazur/University of Gothenburg

The Thwaites Glacier in West Antarctica—about the size of Florida—has been an elephant in the room for scientists trying to make global sea level rise predictions.

This massive ice stream is already in a phase of fast retreat (a "collapse" when viewed on geological timescales) leading to widespread concern about exactly how much, or how fast, it may give up its ice to the ocean.

The potential impact of Thwaites' retreat is spine-chilling: a total loss of the glacier and surrounding icy basins could raise sea level from 3 to 10 feet.

A new study in *Nature Geoscience* led by marine geophysicist Alastair Graham at the University of South Florida's College of Marine Science adds cause for concern. For the first time, scientists mapped in high-resolution a critical area of the seafloor in front of the glacier that gives them a window into how fast Thwaites retreated and moved in the past.

The stunning imagery shows geologic features that are new to science, and also provides a kind of crystal ball to see into Thwaites' future. In people and ice sheets alike, past behavior is key to understanding future behavior.

The team documented more than 160 parallel ridges that were created, like a footprint, as the glacier's leading edge retreated and bobbed up and down with the daily tides.

"It's as if you are looking at a tide gauge on the seafloor," Graham said. "It really blows my mind how beautiful the data are."



THOR scientists Alastair Graham (right) and Robert Larter (left) look on in awe at the crumbling ice face of the Thwaites Glacier margin, from the bridge deck of the R/V Nathaniel B. Palmer. Credit: Frank Nitsche

Beauty aside, what's alarming is that the rate of Thwaites' retreat that scientists have documented more recently are small compared to the fastest rates of change in its past, said Graham.

To understand Thwaites' past retreat, the team analyzed the rib-like formations submerged 700 meters (just under half a mile) beneath the polar ocean and factored in the tidal cycle for the region, as predicted by computer models, to show that one rib must have been formed every

single day.


At some point in the last 200 years, over a duration of less than six months, the front of the glacier lost contact with a seabed ridge and retreated at a rate of more than 2.1 kilometers per year (1.3 miles per year)—twice the rate documented using satellites between 2011 and 2019.

"Our results suggest that pulses of very rapid retreat have occurred at Thwaites Glacier in the last two centuries, and possibly as recently as the mid-20th Century," Graham said.

"Thwaites is really holding on today by its fingernails, and we should expect to see big changes over small timescales in the future—even from one year to the next—once the glacier retreats beyond a shallow ridge in its bed," said marine geophysicist and study co-author Robert Larter from the British Antarctic Survey.

To collect the imagery and supporting geophysical data, the team, which included scientists from the United States, the United Kingdom and Sweden, launched a state-of-the-art orange robotic vehicle loaded with imaging sensors called "Rán" from the R/V Nathaniel B. Palmer during an expedition in 2019.

Rán, operated by scientists at the University of Gothenburg in Sweden, embarked on a 20-hour mission that was as risky as it was serendipitous, Graham said. It mapped an area of the seabed in front of the glacier about the size of Houston—and did so in extreme conditions during an unusual summer notable for its lack of sea ice.



A 3D-rendered view of the multibeam bathymetry (seafloor shape) colored by depth, collected by Rán across a seabed ridge, just in front of Thwaites Ice Shelf. Credit: Alastair Graham/University of South Florida

This allowed scientists to access the glacier front for the first time in history.

"This was a pioneering study of the ocean floor, made possible by recent technological advancements in autonomous ocean mapping and a bold decision by the Wallenberg foundation to invest into this research infrastructure," said Anna Wåhlin, a physical oceanographer from the University of Gothenburg who deployed Rán at Thwaites. "The images Rán collected give us vital insights into the processes happening at the critical junction between the glacier and the ocean today."

"It was truly a once in a lifetime mission," said Graham, who said the team would like to sample the seabed sediments directly so they can more accurately date the ridge-like features.

"But the ice closed in on us pretty quickly and we had to leave before we

could do that on this expedition," he said.

While many questions remain, one thing's for sure: It used to be that scientists thought of the Antarctic ice sheets as sluggish and slow to respond, but that's simply not true, said Graham.

"Just a small kick to Thwaites could lead to a big response," he said.

According to the United Nations, roughly 40% of the human population lives within 60 miles of the coast.



Rán, a Kongsberg HUGIN autonomous underwater vehicle, amongst sea ice in front of Thwaites Glacier, after a 20-hour mission mapping the seafloor. Credit:

Anna Wåhlin/University of Gothenburg

"This study is part of a cross-disciplinary collective effort to understand the Thwaites Glacier system better," said Tom Frazer, dean of the USF College of Marine Science, "and just because it's out of sight, we can't have Thwaites out of mind. This study is an important step forward in providing essential information to inform global planning efforts."

The 2019 expedition was the first in a five-year project dubbed THOR, which stands for Thwaites Offshore Research, and also included team members from a sister project called the Thwaites-Amundsen Regional Survey and Network Integrating Atmosphere-Ice-Ocean Processes, or TARSAN.

More information: Rapid retreat of Thwaites Glacier in the pre-satellite era, *Nature Geoscience* (2022).

doi.org/10.1038/s41561-022-01019-9 ,
www.nature.com/articles/s41561-022-01019-9

Provided by University of South Florida

Citation: New seafloor images off the West Antarctic Ice Sheet upend understanding of Thwaites Glacier retreat (2022, September 5) retrieved 26 June 2024 from <https://phys.org/news/2022-09-seafloor-images-west-antarctic-ice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.